

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-[(*E*)-(4-Ethylphenyl)iminomethyl]-benzene-1,2-diol

 Zeynep Keleşoğlu,^{a*} Orhan Büyükgüngör,^a Çiğdem Albayrak^b and Mustafa Odabaşoğlu^c
^aDepartment of Physics, Ondokuz Mayıs University, TR-55139 Samsun, Turkey,

^bSinop University, Sinop Faculty of Education, Sinop, Turkey, and ^cPamukkale University, Denizli Technical Vocational School, Denizli, Turkey

Correspondence e-mail: zeynep.kelesoglu@omu.edu.tr


Received 21 July 2009; accepted 28 July 2009

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.051; wR factor = 0.148; data-to-parameter ratio = 14.0.

The title compound, $\text{C}_{15}\text{H}_{15}\text{NO}_2$, adopts the enol-imine tautomeric form. The dihedral angle between the two benzene rings is $48.1(1)^\circ$. Intramolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds generate $S(6)$ and $S(5)$ ring motifs, respectively. In the crystal, molecules are linked into centrosymmetric $R_2^2(10)$ dimers via pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds and the dimers may interact through very weak by $\pi-\pi$ interactions [centroid-centroid distance = $4.150(1)$ Å]. The ethyl group is disordered over two orientations, with occupancies of 0.587 (11) and 0.413 (11).

Related literature

For the photochromic and thermochromic properties of Schiff base compounds, see: Elmali *et al.* (1999); Guha *et al.* (2000); Kletski *et al.* (1997); Kownacki *et al.* (1994); Zgierski *et al.* (2000). For Schiff base tautomerism, see: Alarcon *et al.* (1995); Dudek *et al.*, (1966); Salman *et al.* (1991, 1993). For a related structure, see: Özek *et al.* (2009). For graph-set motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{15}\text{NO}_2$	$\alpha = 87.326(6)^\circ$
$M_r = 241.28$	$\beta = 86.397(6)^\circ$
Triclinic, $P\bar{1}$	$\gamma = 69.394(5)^\circ$
$a = 6.1893(4)$ Å	$V = 646.85(8)$ Å ³
$b = 8.7704(6)$ Å	$Z = 2$
$c = 12.7605(9)$ Å	Mo $K\alpha$ radiation

 $\mu = 0.08$ mm⁻¹
 $T = 296$ K

 $0.54 \times 0.41 \times 0.31$ mm

Data collection

Stoe IPDS II diffractometer	8683 measured reflections
Absorption correction: integration (<i>X-RED32</i> ; Stoe & Cie, 2002)	2668 independent reflections
$T_{\min} = 0.966$, $T_{\max} = 0.979$	1896 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.148$	$\Delta\rho_{\text{max}} = 0.24$ e Å ⁻³
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.13$ e Å ⁻³
2668 reflections	
191 parameters	
28 restraints	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.95 (3)	1.72 (3)	2.596 (2)	152 (2)
$\text{O2}-\text{H2}\cdots\text{O1}$	0.88 (3)	2.29 (3)	2.7307 (19)	111 (2)
$\text{O2}-\text{H2}\cdots\text{O1}^{\dagger}$	0.88 (3)	2.06 (3)	2.818 (2)	143 (2)

 Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDS II diffractometer (purchased under grant F.279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2862).

References

- Alarcon, S. H., Olivieri, A. C. & Nordon, A. (1995). *Tetrahedron*, **51**, 4619–4626.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Dudek, G. O. & Dudek, E. P. (1966). *J. Am. Chem. Soc.* **88**, 2407–2412.
- Elmali, A., Kabak, M., Kavlakoglu, E., Elerman, Y. & Durlu, T. N. (1999). *J. Mol. Struct.* **510**, 207–214.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Guha, D., Mandal, A., Koll, A., Filarowski, A. & Mukherjee, S. (2000). *Spectrochim. Acta A*, **56**, 2669–2677.
- Kletski, M., Milov, A., Metelisa, A. & Knyazhansky, M. (1997). *J. Photochem. Photobiol. A*, **110**, 267–270.
- Kownacki, K., Mordzinski, A., Wilbrandt, R. & Grobowska, A. (1994). *Chem. Phys. Lett.* **227**, 270–276.
- Özek, A., Büyükgüngör, O., Albayrak, Ç. & Odabaşoğlu, M. (2009). *Acta Cryst.* **E65**, o791.
- Salman, S. R., Lindon, J. C. & Farrant, R. D. (1991). *Spectrosc. Lett.* **24**, 1071–1078.
- Salman, S. R., Lindon, J. C. & Farrant, R. D. (1993). *Magn. Reson. Chem.* **31**, 991–994.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Zgierski, M. & Grobowska, A. (2000). *J. Chem. Phys.* **113**, 7845–7852.

supporting information

Acta Cryst. (2009). E65, o2055 [doi:10.1107/S1600536809029924]

3-[(*E*)-(4-Ethylphenyl)iminomethyl]benzene-1,2-diol

Zeynep Keleşoğlu, Orhan Büyükgüngör, Çiğdem Albayrak and Mustafa Odabaşoğlu

S1. Comment

There has been considerable interest in some Schiff bases derived from salicylaldehyde and substituted salicylaldehyde because they show thermochromism and photochromism in the solid state (Kownacki *et al.*, 1994). The tautomerism in the Schiff base ligands plays an important role for distinguishing their photochromic (Guha *et al.*, 2000) and thermochromic (Zgierski *et al.*, 2000) characteristics. It has been proposed that molecules showing thermochromism are planar, while those showing photochromism are non-planar (Kletski *et al.*, 1997), both phenomena being associated with a proton transfer (Elmali *et al.*, 1999). Schiff bases derived from the condensation of salicylaldehyde with aniline and substituted aniline, and naphthaldehyde with aniline exists as enol-imine (Dudek *et al.*, 1966), keto-amine (Salman *et al.*, 1991), or enol-imine/keto-amine form (Salman *et al.*, 1993; Alarcon *et al.*, 1995) in all solvents.

The X-ray analysis shows that the title compound prefers an enol-imine tautomeric form, with a strong intramolecular O1—H1···N1 hydrogen bond. This is also confirmed by the C2—O1 [1.361 (2) Å], C7—N1 [1.278 (2) Å], C1—C7 [1.445 (2) Å] and C1—C2 [1.399 (2) Å] bond lengths (Fig. 1). The C2—O1 bond length of 1.361 (2) Å indicates a single-bond character and the C7—N1 bond length of 1.278 (2) Å indicates a high degree of double-bond character. Similar results were observed for (*E*)-4-methoxy-2-[(*o*-tolylimino)methyl]phenol [C—O = 1.357 (2) Å, C=N = 1.286 (2) Å; Özek *et al.*, 2009]. An intramolecular O2—H2···O1 hydrogen bond is also observed. The O—H···N and O—H···O hydrogen bonds generate S(6) and S(5) ring motifs, respectively (Bernstein *et al.*, 1995).

The dihedral angle between benzene rings A(C1-C6) and B(C8-C13) is 48.1 (1)°. The nearly planar S(6) ring C(O1/H1/N1/C1/C2/C7) is oriented with respect to rings A and B at dihedral angles of A/C = 1.89 (42)° and B/C = 46.23 (25)°. It is known that Schiff bases may exhibit thermochromism or photochromism, depending on the planarity or non-planarity of the molecule, respectively. Since the title molecule is non-planar, one can expect photochromic properties in title compound.

In the crystal structure, molecules are linked into centrosymmetric $R_2^2(10)$ dimers via O—H···O hydrogen bonds (Table 2). A very weak π - π interaction occurs between A(C1-C6) rings at (x, y, z) and (1-x, 1-y, 1-z), with a ring centroid-to-centroid distance of 4.150 (1) Å; only atoms C1, C2 and C3 are involved in the interactions as the rings are displaced.

S2. Experimental

Compound (I) was prepared by refluxing a mixture of 2,3-dihydroxy benzaldehyde (0.5 g 0.0036 mol) in ethanol (20 ml) and 4-ethylaniline (0.436 g 0.0036 mol) in ethanol (20 ml). The reaction mixture was stirred for 1 h under reflux. Single crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of a methanol solution (yield 87%, m.p. 378-379 K).

S3. Refinement

The ethyl group is disordered over two orientations, with occupancies of 0.587 (11) and 0.413 (11). The U^{ij} parameters of the disordered atoms were restrained to an approximate isotropic behaviour. The C—C distances involving disordered atoms were restrained to 1.54 (2) Å. The hydroxyl H atoms were located in a difference Fourier map and were refined freely. All other H-atoms were refined using a riding model with $d(\text{C-H}) = 0.93\text{--}0.96$ Å ($U_{\text{iso}} = 1.2U_{\text{eq}}$ of the parent atom) for aromatic and ethyl C atoms and $d(\text{C-H}) = 0.97$ Å ($U_{\text{iso}} = 1.5U_{\text{eq}}$ of the parent atom) for methyl C atoms.




Figure 1

The molecular structure of (I), showing 30% probability displacement ellipsoids and the atom-numbering scheme. Only the major disorder component of the ethyl group is shown. Dashed lines indicate hydrogen bonds.




Figure 2

A packing diagram for (I), showing the formation dimers through O—H...O hydrogen bonds and π - π interactions.

[Symmetry code: (i) $-x, 1 - y, 1 - z$; (ii) $1 - x, 1 - y, 1 - z$]. H atoms not involved in hydrogen bonding (dashed lines) have been omitted for clarity. Cg1 and Cg2 are centroids of the C1-C6 and C8-C13 rings, respectively.

3-[(*E*)-(4-Ethylphenyl)iminomethyl]benzene-1,2-diol

Crystal data

$C_{15}H_{15}NO_2$

$M_r = 241.28$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.1893$ (4) Å

$b = 8.7704$ (6) Å

$c = 12.7605$ (9) Å

$\alpha = 87.326$ (6)°

$\beta = 86.397$ (6)°

$\gamma = 69.394$ (5)°

$V = 646.85$ (8) Å³

$Z = 2$

$F(000) = 256$

$D_x = 1.239$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8683 reflections

$\theta = 1.6$ – 28.0 °

$\mu = 0.08$ mm⁻¹

$T = 296$ K

Prism, red

$0.54 \times 0.41 \times 0.31$ mm

Data collection

Stoe IPDS II

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 6.67 pixels mm⁻¹

rotation method scans

Absorption correction: integration

(*X-RED32*; Stoe & Cie, 2002)

$T_{\min} = 0.966$, $T_{\max} = 0.979$

8683 measured reflections

2668 independent reflections

1896 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$
 $\theta_{\text{max}} = 26.5^\circ$, $\theta_{\text{min}} = 1.6^\circ$
 $h = -7 \rightarrow 7$

$k = -11 \rightarrow 11$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.148$
 $S = 1.03$
 2668 reflections
 191 parameters
 28 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0674P)^2 + 0.1042P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.24 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.13 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $F_c^* = kFc[1 + 0.001x \text{Fc}^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.030 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.5280 (3)	0.5619 (2)	0.34375 (15)	0.0590 (5)	
C2	0.3189 (3)	0.5824 (2)	0.40019 (14)	0.0557 (4)	
C3	0.2107 (3)	0.7228 (2)	0.45769 (15)	0.0583 (5)	
C4	0.3058 (3)	0.8430 (2)	0.45564 (17)	0.0664 (5)	
H4	0.2322	0.9372	0.4932	0.080*	
C5	0.5101 (4)	0.8250 (2)	0.39809 (18)	0.0717 (6)	
H5	0.5719	0.9076	0.3964	0.086*	
C6	0.6212 (4)	0.6859 (2)	0.34379 (17)	0.0700 (5)	
H6	0.7600	0.6736	0.3065	0.084*	
C7	0.6472 (3)	0.4135 (2)	0.28805 (15)	0.0636 (5)	
H7	0.7913	0.3996	0.2554	0.076*	
C8	0.6932 (3)	0.1541 (2)	0.23042 (15)	0.0629 (5)	
C9	0.9261 (4)	0.0745 (3)	0.24457 (18)	0.0752 (6)	
H9	1.0026	0.1169	0.2892	0.090*	
C10	1.0450 (4)	-0.0671 (3)	0.1929 (2)	0.0914 (8)	
H10	1.2010	-0.1202	0.2041	0.110*	
C11	0.9377 (5)	-0.1323 (3)	0.1246 (2)	0.0994 (8)	
C12	0.7038 (5)	-0.0553 (3)	0.1148 (2)	0.0949 (8)	
H12	0.6263	-0.0994	0.0718	0.114*	
C13	0.5816 (4)	0.0857 (3)	0.16706 (18)	0.0789 (6)	

H13	0.4233	0.1347	0.1595	0.095*	
C14A	1.0861 (17)	-0.2683 (8)	0.0432 (7)	0.120 (3)	0.587 (11)
H14A	0.9952	-0.2714	-0.0153	0.144*	0.587 (11)
H14B	1.2235	-0.2474	0.0164	0.144*	0.587 (11)
C15A	1.1473 (16)	-0.4208 (9)	0.1053 (5)	0.143 (3)	0.587 (11)
H15A	1.2155	-0.5112	0.0597	0.214*	0.587 (11)
H15B	1.0107	-0.4296	0.1409	0.214*	0.587 (11)
H15C	1.2558	-0.4216	0.1561	0.214*	0.587 (11)
C14B	1.0391 (18)	-0.3057 (11)	0.0870 (10)	0.116 (4)	0.413 (11)
H14C	1.0239	-0.3787	0.1441	0.139*	0.413 (11)
H14D	0.9493	-0.3172	0.0303	0.139*	0.413 (11)
C15B	1.273 (2)	-0.3543 (17)	0.0517 (11)	0.164 (5)	0.413 (11)
H15D	1.2997	-0.4273	-0.0053	0.246*	0.413 (11)
H15E	1.3686	-0.4085	0.1082	0.246*	0.413 (11)
H15F	1.3090	-0.2601	0.0282	0.246*	0.413 (11)
N1	0.5630 (3)	0.30067 (18)	0.28177 (13)	0.0637 (4)	
O1	0.2152 (2)	0.46864 (15)	0.40254 (11)	0.0659 (4)	
O2	0.0118 (2)	0.74271 (17)	0.51685 (13)	0.0749 (5)	
H1	0.314 (5)	0.386 (3)	0.358 (2)	0.099 (8)*	
H2	-0.027 (5)	0.655 (4)	0.515 (2)	0.111 (9)*	

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0606 (11)	0.0534 (10)	0.0622 (11)	-0.0192 (8)	-0.0047 (9)	0.0012 (8)
C2	0.0577 (10)	0.0463 (9)	0.0641 (11)	-0.0189 (8)	-0.0078 (8)	-0.0009 (7)
C3	0.0546 (10)	0.0491 (9)	0.0695 (12)	-0.0150 (8)	-0.0065 (8)	-0.0049 (8)
C4	0.0681 (12)	0.0502 (10)	0.0818 (13)	-0.0197 (9)	-0.0128 (10)	-0.0079 (9)
C5	0.0766 (13)	0.0580 (11)	0.0896 (15)	-0.0339 (10)	-0.0112 (11)	0.0003 (10)
C6	0.0674 (12)	0.0667 (12)	0.0807 (14)	-0.0301 (10)	-0.0005 (10)	0.0004 (10)
C7	0.0628 (11)	0.0606 (11)	0.0638 (12)	-0.0185 (9)	0.0036 (9)	-0.0008 (9)
C8	0.0712 (12)	0.0564 (10)	0.0585 (11)	-0.0195 (9)	0.0014 (9)	-0.0034 (8)
C9	0.0736 (13)	0.0669 (12)	0.0793 (14)	-0.0156 (10)	-0.0063 (10)	-0.0143 (10)
C10	0.0826 (16)	0.0733 (14)	0.1042 (19)	-0.0078 (12)	-0.0029 (13)	-0.0206 (13)
C11	0.112 (2)	0.0714 (14)	0.104 (2)	-0.0159 (14)	-0.0015 (15)	-0.0307 (13)
C12	0.117 (2)	0.0750 (15)	0.0955 (18)	-0.0321 (15)	-0.0172 (15)	-0.0207 (13)
C13	0.0826 (15)	0.0695 (13)	0.0861 (15)	-0.0265 (11)	-0.0127 (12)	-0.0062 (11)
C14A	0.144 (6)	0.085 (4)	0.129 (5)	-0.042 (4)	0.014 (4)	0.013 (3)
C15A	0.182 (7)	0.098 (5)	0.114 (4)	-0.007 (4)	0.007 (4)	-0.021 (3)
C14B	0.137 (6)	0.059 (5)	0.136 (7)	-0.016 (4)	0.034 (5)	-0.044 (5)
C15B	0.156 (8)	0.135 (7)	0.185 (9)	-0.036 (6)	0.038 (7)	-0.029 (6)
N1	0.0668 (10)	0.0559 (9)	0.0655 (10)	-0.0181 (7)	0.0016 (7)	-0.0054 (7)
O1	0.0642 (8)	0.0517 (7)	0.0840 (10)	-0.0234 (6)	0.0090 (7)	-0.0149 (6)
O2	0.0646 (9)	0.0587 (8)	0.1033 (12)	-0.0236 (7)	0.0119 (7)	-0.0265 (7)

Geometric parameters (Å, °)

C1—C2	1.399 (3)	C11—C12	1.376 (4)
C1—C6	1.399 (3)	C11—C14B	1.514 (7)
C1—C7	1.445 (3)	C11—C14A	1.603 (8)
C2—O1	1.361 (2)	C12—C13	1.379 (3)
C2—C3	1.395 (2)	C12—H12	0.93
C3—O2	1.364 (2)	C13—H13	0.93
C3—C4	1.375 (3)	C14A—C15A	1.464 (11)
C4—C5	1.385 (3)	C14A—H14A	0.97
C4—H4	0.93	C14A—H14B	0.97
C5—C6	1.367 (3)	C15A—H15A	0.96
C5—H5	0.93	C15A—H15B	0.96
C6—H6	0.93	C15A—H15C	0.96
C7—N1	1.278 (2)	C14B—C15B	1.405 (15)
C7—H7	0.93	C14B—H14C	0.97
C8—C13	1.378 (3)	C14B—H14D	0.97
C8—C9	1.382 (3)	C15B—H15D	0.96
C8—N1	1.419 (2)	C15B—H15E	0.96
C9—C10	1.375 (3)	C15B—H15F	0.96
C9—H9	0.93	O1—H1	0.95 (3)
C10—C11	1.383 (4)	O2—H2	0.88 (3)
C10—H10	0.93		
C2—C1—C6	118.93 (17)	C10—C11—C14A	121.0 (4)
C2—C1—C7	120.31 (17)	C11—C12—C13	121.7 (2)
C6—C1—C7	120.76 (18)	C11—C12—H12	119.2
O1—C2—C3	117.70 (17)	C13—C12—H12	119.2
O1—C2—C1	122.45 (16)	C8—C13—C12	120.2 (2)
C3—C2—C1	119.85 (16)	C8—C13—H13	119.9
O2—C3—C4	119.15 (17)	C12—C13—H13	119.9
O2—C3—C2	121.02 (16)	C15A—C14A—C11	104.1 (6)
C4—C3—C2	119.83 (18)	C15A—C14A—H14A	110.9
C3—C4—C5	120.55 (18)	C11—C14A—H14A	110.9
C3—C4—H4	119.7	C15A—C14A—H14B	110.9
C5—C4—H4	119.7	C11—C14A—H14B	110.9
C6—C5—C4	120.14 (19)	H14A—C14A—H14B	109.0
C6—C5—H5	119.9	C14A—C15A—H15A	109.5
C4—C5—H5	119.9	C14A—C15A—H15B	109.5
C5—C6—C1	120.7 (2)	H15A—C15A—H15B	109.5
C5—C6—H6	119.7	C14A—C15A—H15C	109.5
C1—C6—H6	119.7	H15A—C15A—H15C	109.5
N1—C7—C1	122.76 (18)	H15B—C15A—H15C	109.5
N1—C7—H7	118.6	C15B—C14B—C11	114.5 (9)
C1—C7—H7	118.6	C15B—C14B—H14C	108.6
C13—C8—C9	118.79 (18)	C11—C14B—H14C	108.6
C13—C8—N1	118.77 (19)	C15B—C14B—H14D	108.6
C9—C8—N1	122.40 (18)	C11—C14B—H14D	108.6

C10—C9—C8	120.3 (2)	H14C—C14B—H14D	107.6
C10—C9—H9	119.9	C14B—C15B—H15D	109.5
C8—C9—H9	119.9	C14B—C15B—H15E	109.5
C9—C10—C11	121.5 (2)	H15D—C15B—H15E	109.5
C9—C10—H10	119.3	C14B—C15B—H15F	109.5
C11—C10—H10	119.3	H15D—C15B—H15F	109.5
C12—C11—C10	117.4 (2)	H15E—C15B—H15F	109.5
C12—C11—C14B	116.2 (5)	C7—N1—C8	120.19 (17)
C10—C11—C14B	123.6 (5)	C2—O1—H1	103.8 (15)
C12—C11—C14A	120.4 (4)	C3—O2—H2	111.0 (18)
C6—C1—C2—O1	178.84 (17)	C9—C10—C11—C12	-3.7 (4)
C7—C1—C2—O1	-2.3 (3)	C9—C10—C11—C14B	-164.1 (6)
C6—C1—C2—C3	-1.9 (3)	C9—C10—C11—C14A	164.1 (4)
C7—C1—C2—C3	177.02 (17)	C10—C11—C12—C13	2.8 (4)
O1—C2—C3—O2	2.1 (3)	C14B—C11—C12—C13	164.7 (6)
C1—C2—C3—O2	-177.26 (17)	C14A—C11—C12—C13	-165.0 (4)
O1—C2—C3—C4	-178.37 (17)	C9—C8—C13—C12	-3.3 (3)
C1—C2—C3—C4	2.3 (3)	N1—C8—C13—C12	179.0 (2)
O2—C3—C4—C5	178.65 (18)	C11—C12—C13—C8	0.6 (4)
C2—C3—C4—C5	-0.9 (3)	C12—C11—C14A—C15A	-109.9 (7)
C3—C4—C5—C6	-0.9 (3)	C10—C11—C14A—C15A	82.7 (8)
C4—C5—C6—C1	1.4 (3)	C14B—C11—C14A—C15A	-21.5 (12)
C2—C1—C6—C5	0.0 (3)	C12—C11—C14B—C15B	151.7 (11)
C7—C1—C6—C5	-178.83 (18)	C10—C11—C14B—C15B	-47.7 (16)
C2—C1—C7—N1	4.6 (3)	C14A—C11—C14B—C15B	45.5 (13)
C6—C1—C7—N1	-176.53 (19)	C1—C7—N1—C8	-176.85 (17)
C13—C8—C9—C10	2.5 (3)	C13—C8—N1—C7	-139.4 (2)
N1—C8—C9—C10	-179.9 (2)	C9—C8—N1—C7	43.0 (3)
C8—C9—C10—C11	1.1 (4)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots N1	0.95 (3)	1.72 (3)	2.596 (2)	152 (2)
O2—H2 \cdots O1	0.88 (3)	2.29 (3)	2.7307 (19)	111 (2)
O2—H2 \cdots O1 ⁱ	0.88 (3)	2.06 (3)	2.818 (2)	143 (2)

Symmetry code: (i) $-x, -y+1, -z+1$.