



MINISTRY OF ENVIRONMENT AND URBANIZATION  
General Directorate of the Protection of  
Natural Assets



MINISTRY OF FORESTRY AND WATER AFFAIRS  
General Directorate of Nature Conservation  
and National Parks



MUĞLA GOVERNORSHIP



# CARETTA CARETTA

FOR SUSTAINABLE SEAS PLEASE HELP PROJECT THE CARETTA CARETTA










DEKAMER-TURKEY

MONITORING AND CONSERVATION STUDIES OF  
SEA TURTLES (*Caretta caretta*) DURING THE 2016 NESTING  
SEASON ON MUĞLA SEA TURTLE NESTING BEACHES

## ACKNOWLEDGMENTS

We would like to thank to:

-  The Ministry of Environment and Urbanization (General Manager and Staff of General Director of The General Directorate of the Protection of Natural Assets),
-  The Ministry of Forestry and Water Affairs (General Managers and Staff of Provincial Directorate of Environment and Urbanization for their supports),
-  Governor of Muğla Province Mr. Amir Çiçek and Governor of Ortaca District Mr. Fatih Ürkmezer,
-  For Koycegiz-Dalyan Region: Directorate of Commercial Enterprise of Muğla Sıtkı Koçman University and Their Staff and the Rector of Muğla Sıtkı Koçman University,
-  Rector and Directors of Pamukkale University for invaluable support to DEKAMER,
-  Directors and personnel of Regional Directorate of Muğla Environment and Urbanization for supports during our field work,
-  And personnel and volunteers of Pamukkale University Sea Turtle Research, Rescue and Rehabilitation Center – DEKAMER,

for their supports during the course of the conservation and monitoring projects in Koycegiz-Dalyan Specially Protected Area, Fethiye Specially Protected Area and Dalaman-Sarıgerme Sea Turtle Nesting Beach during 2016 sea turtle breeding season.

**Prof. Dr. Yakup KASKA**

*DEKAMER Manager / Project Coordinator*



## I. SEA TURTLES

There are 8 species of sea turtles are living in the world (Lutz & Musick, 1997).

Table 1: Sea turtle species of the world

Species	English Name
<i>Caretta caretta</i>	Loggerhead Turtle
<i>Chelonia mydas</i>	Green Turtle
<i>Chelonia agassizi</i>	Black Turtle
<i>Dermochelys coriacea</i>	Leatherback Turtle
<i>Eretmochelys imbricata</i>	Hawksbill Turtle
<i>Lepidochelys olivacea</i>	Olive Ridley
<i>Lepidochelys kempii</i>	Kemp's Ridley
<i>Natator depressus</i>	Flatback Turtle


The nesting beaches of sea turtles, which have managed to survive since dinosaurs have dominated the earth, are slowly disappearing. The main reason for this disappearance is the increasing rate of human activities. The injuries and deaths of turtles and plundering of their nests in large numbers have remained a major reason for the decline of populations. Another reason of decline in sea turtle populations has been due to random deaths in uncontrolled hunting and fishing activities for personal or commercial purposes (Lutz & Musick, 1997).

The survivorship of sea turtle populations depends primarily on the conservation of the their nesting beaches they use for reproduction and maintain the mating, feeding, wintering and migratory habitats in their natural ecosystems. Therefore, the beaches in Turkey are of great importance for the Mediterranean population (Başoğlu & Baran, 1982; Baran & Kasperek, 1989; Baran, 1990; Baran et.al., 1992).

## NESTING SPECIES IN THE MEDITERRANEAN



## NON-NESTING SPECIES IN THE MEDITERRANEAN



## SPECIES NOT RECORDED IN THE MEDITERRANEAN




Figure 1: Sea turtle species



## SOME INTERESTING CHARACTERISTICS OF SEA TURTLES

1

Sea turtles show natal homing instincts that bring them back to beaches where they were born to nest.



2

Females nest 3-5 times in every 2-3 years. Interval between each nest is about 15 days.

3

Loggerhead (*Caretta caretta*) turtles build their nests at about 50-60 cm depth on the beach, green (*Chelonia mydas*) turtles at about 90-100 cm depth



4

Clutches of eggs are between 50-100 and each egg is about the size of a ping-pong ball. The incubation period is between 45-65 days. Sexual maturity takes 25-30 years in sea turtles. Only 3-5 of every thousand hatchlings can survive until full maturity.

5

The sex of sea turtles is 'temperature dependent'; higher temperatures (32°C) produce females and lower temperatures (26°C) males.



6

*Caretta caretta* is carnivorous and can dive up to 200 meters sea depth while *Chelonia mydas* is herbivorous and can swim at a depth of 20-50 meters. Sea turtles are 'ecotransformers'. They eat and live in water but move to the land and deposit their eggs.

7

Sea turtles don't have teeth, but they have very strong jaws and saw-shaped palate.



8

They tear and crush food and thereby create small particles of food for other animals.

9

Some sea turtles feed on jellyfishes. They sometimes eat plastic bags instead of jellyfish by mistake, they can suffocate and die.



10

Newly hatched turtles orientate towards reflection of moon and reach to the sea, but they become disorientated when they see any light other than from the sea.

## I. I. Life Cycle of Mediterranean Sea Turtles

Sea turtles maintain a very short period of their lives dependent on the land. Sea turtles mate in the calm waters in the regions close to the nesting beaches in April – May. Sea turtles come to the beach where they borne and lay their eggs between May and August.

The eggs are about the size of a ping pong ball and average 70 eggs in a clutch. The depths of the nests are about 50 cm for loggerhead turtles and relatively deeper for green turtles. The female turtle returns to the sea after leaving her eggs to the beach.

After mating near the nesting beach, the males migrate again to the foraging area. The females return to the foraging and wintering areas after laying their eggs. The offspring are hatched and emerged from the egg at the end of the incubation period of about 2 months. The young offspring travel towards the sea and spends 25–30 years for sexual maturation to come back to the beach where they born (Figure 2).




Figure 2: Life cycle of sea turtles



Sea turtles mate in April and May in shallow waters close to their nesting sites.



Sea turtles lay their eggs between May and August. The size of the egg is similar to size of a ping-pong ball. There are approximately 70 eggs in a nest. The depth of the nests is around 50cm.



Hatchlings emerge from the nests after 2 months of incubation period.



Hatchlings, when they reach the sea, spends around 25 years in the sea and never comes to the beach. Only females come to lay their eggs when they get sexually mature at the age of 25 years.

The number of sea turtles in the world is gradually decreasing. Over the last 500 years, the consumption of sea turtles' meats, eggs, shells, oil, and skin has led these species in danger of extinction. Thousands of sea turtles are drowned each year due to shrimp nets, trawl nets and longlines (Ripple, 1996). Especially the translucent plastic debris are resembled to the jellyfish and being eaten by sea turtles causes deaths or injuries. In addition, collision with marine vehicles cause deaths of mature and juvenile sea turtles.

One of the problems that adult sea turtles have experienced on land is the occupation of the nesting beaches in recent years and as a result, the generations of sea turtles, which have been living for millions of years, are getting more and more threatened every day. In this way, the usage plans of the coastal habitats which are used as nesting beaches by sea turtles have changed, and as a result, they have become endangered (Ozaner, 1991).

In recent years, the predation of nests by predator animals such as foxes, badgers, dogs and crabs has increased considerably and the measures taken have not been able to respond in a good level. In addition, due to the effect of the artificial lights behind the beaches, the hatchlings are disoriented by these lights and lost their ways and die without finding the sea (Witherington & Bjorndal, 1991). The nests close to the sea can face the danger of inundation, and as a result, the hatchlings die before they complete their embryonic development. Another factor affecting turtle nests is the activity of insects that invade nests. Marine pollution also affects juvenile turtles as adult turtles. The predation usually observed on sea turtle eggs and their hatchlings. Foxes, badgers and crabs are main predators in Turkish beaches. Fox and badger predation directly affect the nest and damage the eggs.

In nesting beaches, another factor damaging sea turtle eggs and hatchlings is humans. Digging sand may damage the eggs and also the holes dug by people becomes a trap for hatchlings. When they fall into these holes they can't reach to the sea and eventually die or predated by other animals.

Another factor is use of sun umbrellas (parasols) use on the nesting beaches. When eggs are found in areas where the parasols are used, the eggs may be damaged directly. In addition, the use of fire at night and heavy vehicles on the beach are also negative factors that caused by people.

## I. 2. Sea Turtles in The Mediterranean and Turkey

Five sea turtle species (*C. caretta*, *C. mydas*, *E. imbricata*, *D. coriacea* and *L. kempii*) were reported in previous studies (Başoğlu, 1973; Groombridge, 1990). Of these, only *C. caretta* and *C. mydas* are nesting in Turkey (Baran, 1990; Baran et al., 1992; Baran & Kasperek, 1989; Başoğlu, 1973; Canbolat, 1991; Geldiay, 1983, 1984; Geldiay & Koray, 1982; Hathaway, 1972). In addition, there are a few stranding reports about *D. coriacea* but all specimens found dead (Baran et al., 1998; Taşkınak et al., 1998; Sönmez et al., 2008).

The number of adult female loggerheads in the world are estimated as 60.000. Only 2.000 of them are living in the Mediterranean and 450-900 females are nesting in Turkey. Adult female green turtle population is estimated to be 200.000 in the world while only 500 of them are living in the Mediterranean and 230-380 of them are nesting in Turkey.

## I. 3. Sea Turtle Studies in Turkey and Muğla Beaches

Approximately 30% to 50% of loggerheads and 70% to 80% of green turtles that are living in the Mediterranean are nesting on the coasts of Turkey. There are 21 main sea turtle nesting beaches in Turkey. These beaches from west to east are; Ekincik, Dalyan, Dalaman-Sarıgerme, Fethiye, Patara, Kale-Demre, Finike-Kumluca, Çıralı-Olimpos, Tekirova, Belek, Kızılot, Demirtaş, Gazipaşa, Anamur, Göksu Deltası, Alata, Davultepe, Kazanlı, Akyatan-Karataş, Sugözü-Yumurtalık and Samandağ (Figure 3). The annual number of nests on these beaches are given in Table 2.




Figure 3 : Main sea turtle nesting beaches in Turkey.  
(Highlighted nesting beach names are indicating main nesting beaches in Turkey)



Table 2: The number of nest range of *C. caretta* and *C. mydas* on nesting beaches.

Beach Number	Beach Name	Length of the beach (km)	The range of nests numbers	
			<i>C. caretta</i>	<i>C. mydas</i>
1	Ekincik	1	9-12	
2	Dalyan	4.7	57-658	
3	Dalaman-Sarıgerme	10.4	69-112	
4	Fethiye	8.3	72-191	
5	Patara	14	35-127	2-2
6	Kale-Demre	8.5	39-109	
7	Finike-Kumluca	21	75-305	0-7
8	Çıralı-Olimpos	3.2	23-139	
9	Tekirova	3.7	4-23	
10	Belek	29.3	68-1900	2-8
11	Kızılot	16.1	50-270	0-3
12	Demirtaş	7.8	41-137	
13	Gazipaşa	6.8	14-53	
14	Anamur	12.2	146-1240	1-3
15	Göksu Deltası	25.6	36-254	3-20
16	Alata	3	16-32	20-356
17	Daultepe	1.8	2-11	86-126
18	Kazanlı	4.5	2-26	73-856
19	Akyatan-Karataş	22	3-31	108-735
20	Sugözü-Yumurtalık	9.4	1-2	126-213
21	Samandağ	14.2	7-20	40-1172

The number of nests obtained from the conservation and monitoring studies since the beginning of 90's were given for Dalyan, Dalaman-Sarıgerme and Fethiye beaches in Muğla. In general, the average of annual number of nests on these three main nesting beaches in Muğla since 2002 is 511 (Figure 4).






Figure 4 : Annual nest numbers on three nesting beaches of Muğla (Geldiay et al. (1982); Canbolat (1991); Erk'akan (1993); Baran et al. 1992, Canbolat (1997, 2001c, 2002, 2003, 2004, 2006, 2007); Yerli and Demirayak, (1996); Baran et al. (1996); Ilgaz and Baran (2001); Yerli and Canbolat (1998b), Türkozan and Vilmaz (2008); Kaska et al. (2008); Durmuş and Güçlü (2009); Kaska et al. (2010, 2011, 2012, 2013, 2014, 2015, 2016) Fethiye - (Türkozan & Baran, 1996; Baran & Türkozan, 1996; Türkozan, 2000, 2006; Yerli & Canbolat, 1998; Ilgaz et al., 2007; Canbolat, 2006, 2007; Takva, 2008, 2009; Canbolat et al., 2010; Başkale et al., 2012, 2013, 2016; Hoş et al., 2014; Yerli et al., 2015; Kaska et al., 2016)



## 2. MATERIAL AND METHODS - 2. I. Study Areas




Figure5: Satellite images of three nesting beaches in Muğla

## 2.2. Methodology

The methodology for determining and recording nests was used according to the previous published studies (Kaska et al., 1998; Başkale & Kaska, 2005; Türkozan & Kaska, 2010). A few photographs showing our studies are given below (Figure 6–10).



Figure 6: Some pictures from field studies



Figure 7: Examples of caging of nests (grid cages against fox predation, on the left, and prisma cages for people attention, on the right)




Figure 8 : Measurements of turtle carapace





Figure 9: Some pictures from field studies



Figure 10: Process of picking up, carrying, measuring of the injured or dead sea turtles

## 3. RESULTS

### 3.1. Results of Sea Turtle Activities

All sea turtle nests spotted on Muğla beaches were belong to loggerhead turtle. A total of 658 nests found on Dalyan beach, while 85 nests on Dalaman-Sarıgerme beach and 112 nests on Fethiye beach. The number of nests found on Dalyan beach has reached the highest number in 2016. GPS coordinates of all nests were taken, the distance from the sea of all nests were also measured and recorded. Nest density on these beaches are; 146.2 nest/km on Dalyan beach, 8.2 nests/km on Dalaman-Sarıgerme beach and 13.5 nests/km on Fethiye beach. The total nest density on these three beaches is 36.54 nests/km in 23.4 km long study area.

A total of 698 nests (81.6%) were screened with grid and prism cages against predation and possible human damages. Of the total nests, a hundred of them (11.7%) were fully predated while 78 of them (9.1%) were half-predated. A total of 677 nests (79.2%) were successfully (without predation) protected.

During 2016 breeding season, a total of 62522 eggs were laid and 42746 hatchlings hatched from these eggs and of these hatchlings, 38193 (89.3%) of them reached to the sea. After hatchling emergences, all nests were uncovered for control to gather data about the nests by the research team (Figure 11). During these controls, a total of 4083 unhatched eggs were identified as unfertilized, 6230 eggs were found with dead embryos in the egg shell. Remaining 9453 eggs were predated by different predators.



Figure 11: Excavation of a nest after hatching








### 3.2. Temporal and Spatial Distribution of Turtle Activities

Evaluating the study results under subsection scale for all three beaches will give more detailed information about turtle activities. Nest number were given under subsections for all three beaches.

In 2016 breeding season, the earliest known sea turtle nesting record for the entire Mediterranean until now was observed on Dalyan Beach on April 25th. The number of nesting activities on Dalyan beach was higher than previous nesting seasons in April and May (19,9%). In general, it is seen that the majority of the nests were made in June. According to this results, the number of nests were six in April (only on Dalyan beach), 164 in May, 459 in June, 221 in July and 5 in August (only on Dalyan beach).




The majority of the nests were made in 10 to 20 m zone from the sea for Dalyan and Fethiye beaches. Nest density is increasing after 20 m from the sea for Dalaman-Sarıgerme. This result shows how important the beach structure is in nest-site selection for the same species' individuals and the importance of making a separate assessment for each beach.

## The Number of Nests and Non-nesting Emergences on the Nesting Beaches of Muğla








## Temporal Distribution of Nests and Non-nesting Emergences on the Nesting Beaches of Muğla



## Nest and Non-nesting Emergence Distributions According to the Distance from the Sea on the Nesting Beaches of Muğla



## The Maps of Nest Locations According to the GPS Coordinates



Sea turtles tend to make their nests in areas that the beach topography allows the highest hatching success. Therefore, the length of the wet and semi-wet area where the sand is washed by the waves is a very important parameter for determining the safe nesting zone. In 2016 breeding season, the sum of the length of the wet sand (WSL) and the length of the semi-wet area that has inundation risk was measured as 8.6 m for Dalyan, 5.2 m for Dalaman-Sarigerme and 5.0 m for Fethiye beaches. The mean distance from the sea of the nests was 15.5m for Dalyan, 39.6 m for Dalaman-Sarigerme and 19.0 m for Fethiye. This is another indicator showing the importance of separate evaluation for each beach.

### 3.3. Results of Tagging and Morphometric Measurements

During night patrols, a total of 165 turtles were encountered on the beaches of Muğla. Of these turtles, 127 of them were tagged during field studies (Dalyan 119; Fethiye 8). Remaining 38 turtles were recaptures from previous years (Dalyan 37; Fethiye 1) (Figure 12).





Figure 12 : Measuring a loggerhead turtle carapace

### 3. 4. Results of Nest Protection and Nest Success



A total of 2.027 sea turtle activities were observed on three nesting beaches in Muğla and 855 of them resulted with nest. Of these nests, 658 of them were determined on Dalyan beach while 85 on Dalaman-Sarıgerme beach and 112 on Fethiye beach. Of these nests, 677 of them were protected. A total of 62.522 eggs were deposited to these nests and 42.746 hatchling were produced on three nesting beaches in Muğla. The information about the nests are given below.

#### Information about Nest Protection (left figures) and Fate of Eggs (right)



##### Fethiye Beach



##### Dalaman - Sarıgerme Beach



##### Dalyan Beach



### 3.5. Sea Turtle Rescue and Rehabilitation

During 2016, a total of 41 sea turtles were admitted to DEKAMER (Sea Turtle Research, Rescue and Rehabilitation Center). Of these, 16 turtles were released back to the sea after successful treatment. Rehabilitation process of 9 turtles are still on going. The number of annual injured turtles admitted to DEKAMER and the cause of injuries are given below (Figure 13-14).




Figure 13: The number of annual injured turtles admitted to DEKAMER




Figure 14: Cause of Injuries of the Sea Turtles Admitted to DEKAMER



# Treatment and Operation Process of Injured Turtles at DEKAMER



## Routine Tank Cleaning Process by Volunteers and Testing Turtles with Alive Preys






## Observing Turtles in the Diving Tanks after Treatment



## RELEASING TURTLES AFTER TREATMENT AND TRACKING WITH SATELLITE TRANSMITTERS





#### 4. PUBLIC AWARENESS STUDIES

Most intensive tourism activities are seen in Dalyan and Fethiye among the nesting beaches of Muğla. Locals, foreign tourists, tourism enterprises in the region and NGO's were informed by project teams and volunteers during the studies.

Public awareness activities on the Dalyan Beach have been made directly to the visitors of DEKAMER and reached to approximately 50,000 people in 2016. Questions about protection sea turtles and of injured turtles have been answered by the DEKAMER staff and volunteers. In addition, the signboards in the center have been renovated and visuals like brochures and booklets have been distributed to visitors. In addition to this, an introduction documentary film about DEKAMER has been shown continuously in the center.

In Fethiye Beach, awareness studies were carried out by distributing hand brochures to the local and foreign beach users in order to protect sea turtles and new warning signs were placed. In addition to this, presentations about sea turtles and biodiversity were made in order to inform the guests and to train the managers and staff of the tourism facilities in Fethiye.

Between June and September, about 100 visitors were informed by the project team and volunteers every evening between 21: 00-23: 30 hours in the information office on Calis Beach. Over 12,000 local and foreign tourists have been informed as a result of public awareness studies during 2016 in Fethiye. A total of 4,000 brochures were also distributed to local and foreign tourists at the information office and during the hotel seminars that held in the hotels.

In addition to this, we interviewed the managers of the tourism facilities and non-governmental organizations in the region and exchanged ideas for the solutions of problems on the beach in order to protect the sea turtle nests.



## INFORMATION TABLES AND INFORMATION SIGNS ON PRISM CAGES ON THE NESTING BEACHES



WE CAN PROTECT SEA TURTLES IF ONLY THE NEW GENERATION LOVES THEM



PUBLIC AWARENESS STUDIES





## PHOTOGRAPHS OF VOLUNTEERS AND THE PROJECT TEAM



## 5. EVALUATION, RESULTS AND RECOMMENDATIONS

### 5. 1. Evaluation the Results of Sea Turtle Monitoring and Conservation Studies

Dalyan, Dalaman-Sarıgerme and Fethiye beaches are the most intense areas among the sea turtle nesting areas in Muğla. A total of 855 nests were made in 2016 in these beaches in Muğla, of these 658 nests in Dalyan was the highest number of nests ever reached. When we look at the results of 24 years of conservation studies for Fethiye, it is seen that the average number of nests is 102, and this year it has reached to above this average with 112 nests. In Dalaman-Sarıgerme, the number of nests has been reached in the average of previous years. The increasing annual nest numbers, especially on Dalyan Beach, can be seen as a result of conservation studies that have been applied for many years but it can be called early to reach a final judgment. Further studies for finding out the other causes of this increase in nest numbers has a great importance in this sense.

While the nest success shows similar characteristics to the previous years for all the beaches, the number of hatchlings reached to the sea has been kept at the same success level achieved in recent years through the intensive conservation efforts on predation in Dalyan. The fox predation was higher in comparison with previous years in Fethiye in 2016 season. Therefore, priority was given to nest caging efforts. In Dalaman-Sarıgerme, it can be said that predation continues as an important problem which reduces the nest success.

Spatial distribution of the nests shows a big variation between nesting zones and each beach's subsections among three different nesting beaches. When the nest densities are considered

according to the distances from the sea, the closest nests are made in Dalyan while the farthest nests are made in Dalaman-Sarıgerme. Thus, the conservation measures should be taken by evaluating each nesting beach separately and it should be avoided taking specific conservation measure for all beaches.


These results are important for the regulation of conservation measures and indicate that conservation & monitoring studies should be continued every year.






## The Annual Number of Eggs and Hatching Success in Muğla Nesting Beaches

Fethiye Beach



Dalaman- Sarigerme Beach



Dalyan Beach



## 5. 2. Suggestions for Nesting Beaches of Muğla

Muğla is one of the most important tourism regions in Turkey. Therefore, the beach and marine area usage is at very high level. Considering that the sea turtles are actively using the coasts of Muğla during summer as breeding and foraging areas, human and sea turtles are facing in many situations. For this reason, conservation and monitoring studies should be continued in all regions and high attention should be given to public information activities. Suggestions for solutions to the problems encountered during the conservation and monitoring studies are given below for each region.

### Correct Light Usage at Nesting Beaches



#### “ SUGGESTION

THE ARTIFICIAL LIGHT SOURCES BEHIND THE NESTING BEACHES IN FETHIYE, PARTICULARLY IN THE HOTELS REGION AND WALKWAYS WHERE INTENSIVE PEOPLE USAGE IS SEEN SHOULD BE CHANGED WITH THE PROPOSED LAMB MODEL GIVEN IN THE FIGURES. LOW SODIUM PRESSURIZED RED LIGHT LAMBS SHOULD BE USED AS LIGHT SOURCE.



### Vehicle Use on the Beach



#### “ SUGGESTION

ALL VEHICLE USAGE ON THE NESTING BEACHES SHOULD BE FORBIDDEN BETWEEN MAY AND SEPTEMBER. ALL KIND OF WORKS THAT REQUIRES VEHICLE USE SHOULD BE COMPLETED BEFORE APRIL.

## Use of Sunbeds on the Nesting Beaches



### “ SUGGESTION

TOURISM FACILITIES ON THE BEACH SHOULD FOLLOW THE SUNBED USAGE REGULATIONS. FACILITIES THAT DO NOT COMPLY WITH THE RULES SHOULD BE PENALIZED BY THE AUTHORITIES. ALSO, LOCALS AND TOURISTS ARE PUTTING THEIR UMBRELLAS. AT LEAST TWO PERSONNEL SHOULD BE CHARGED TO CONTROL THE REGULATIONS TO PREVENT ANY POSSIBLE NEGATIVE ACTIVITIES ON THE BEACH. IN ADDITION, PLACEMENT OF ANY CARPET, WOODEN PLATFORMS ETC. ON THE NESTING BEACHES SHOULD BE PREVENTED.



## Speed Boat Use in the Sea



### “ SUGGESTION

SPEED BOAT USE IN THE SEA SPEED LIMITS SHOULD BE APPLIED TO PREVENT SEA TURTLES FROM BEING DAMAGED FROM SPEED BOATS, AND APPROPRIATE AREAS FOR PERFORMING THESE ACTIVITIES SHOULD BE DETERMINED. NIGHT ACTIVITIES WITH SPEED BOATS SHOULD BE FORBIDDEN. LOCAL TOURISM ENTERPRISES AND FISHERMEN WERE ENCOURAGED TO USE THEIR MARINE VEHICLES IN APPROPRIATE AREAS IN THE LEGAL SPEED LIMITS ACCORDING TO THE LEGISLATIONS.



## Litters on the Beaches



### “ SUGGESTION

LITTERS ON THE BEACH SHOULD BE CLEANED PERIODICALLY AND WARNING SIGNS AND TRASHCANS SHOULD BE PLACED IN APPROPRIATE AREAS ON THE BEACH. USAGE OF PLASTIC BAGS SHOULD BE LIMITED AND PEOPLE SHOULD BE INFORMED ABOUT PLASTIC BAG USAGE.



YOU SEE THE DIFFERENCE.  
**A TURTLE DOES NOT.**



**MEDASSET**  
Mediterranean Association  
to Save the Sea Turtles

[www.medasset.org](http://www.medasset.org)



[www.dekamer.org.tr](http://www.dekamer.org.tr)

## Fox and Dog Predation on Sea Turtle Nests

For more effective conservation applications, nests that under risk of inundation (identified in Section 3.2) or in the areas with high predation, the following measures should be applied:

- Nests that under risk of inundation should be altered to safe areas
- All nests should be screened against predation with grid 1 m wide cages
- Cages should be strengthened with 4 metal stakes on their corners
- Using of 25 cm side-cages around the grid cages should be continued
- The conservation studies has to start on April 15, taking into account the early start of the nesting season



### “ SUGGESTION

SCREENING OF THE NESTS WITH GRID CAGES AGAINST FOX AND DOG PREDATION SHOULD BE CONTINUED AND THE NESTS AT HIGH INUNDATION RISK SHOULD BE ALTERED TO SAFER PLACES BY THE EXPERTS.





## Shower Waters Running Toward the Beach



### SUGGESTION

THE WATER RUNNING FROM THE SHOWERS ARE WETTING THE BEACH AND THREATENING THE SEA TURTLE NESTS. THESE SHOWERS SHOULD BE MOVED ACROSS THE ROAD OR WATER OUTLETS SHOULD BE MADE.

## Feeding Sea Turtles by Hand



### SUGGESTION

BLUE CRAB, ONE OF THE FOOD SOURCES OF LOGGERHEAD TURTLES ARE BEING CAUGHT AT NIGHT IN THE LAGOON BEHIND THE DALYAN BEACH. THIS IS CAUSING LIGHT POLLUTION ON THE BEACH. ALSO, LOCALS ARE USING THESE CRABS FOR TOURISTIC SHOWS TO FEED TURTLES BY HAND. SOME TOURISM AGENCIES ARE PROMISES TO SHOW SEA TURTLES IN THEIR PROGRAM AND FEEDING SEA TURTLES BY HAND REGULARLY. THIS IS AFFECTING TURTLES NEGATIVELY. TURTLES OWN THE AREA AND CHANGE THEIR BEHAVIOR. SOME TURTLES STOP MIGRATING TO THEIR WINTERING AREAS. IN SOME CASES, TURTLES ARE FED BY CHICKEN SKIN INSTEAD OF CRAB OR FISH WHICH CHANGE THEIR DIET AND AFFECTS THEM NEGATIVELY. ALSO, THE CHANGE IN THEIR FEEDING BEHAVIOR AND EXPOSURE TO CONSTANT STRESS IN A CROWDED AREA IS LEADING THEM TO SECONDARY PROBLEMS. IN SOME CASES, TURTLES CAN BITE PEOPLE FROM THEIR LEGS. THUS, FEEDING TURTLES BY HAND SHOULD BE FORBIDDEN AND TOURISTS SHOULD BE INFORMED ABOUT THE NEGATIVE EFFECTS OF THESE ACTIVITIES. EVENTHOUGH THE HAND FEEDING ACTIVITIES WERE BANNED AT THE END OF THE SEASON, SOME TURTLES CONTINUED TO BITE PEOPLE AFTER THE ALTERATION OF THEIR BEHAVIOR.



## Rehabilitation of Sea Turtles in Sea-Pens



### “ SUGGESTION

TWO SEA-PENS WERE PLACED 1 MILE FROM THE BEACH TO OBSERVE REHABILITATED SEA TURTLES DIVING BEHAVIOR AND ADAPTATION TO NATURAL ENVIRONMENT BEFORE RELEASE. THESE SEA-PENS ARE 8 TO 10 M DEEP AND THE DIAMETERS ARE 9 AND 15 M. THESE SEA-PENS CAN ALSO BE USED FOR REHABILITATION OF BITING TURTLES AFTER ALTERED BEHAVIOR AS A RESULT OF HAND FEEDING ACTIVITIES. IN THESE SEA-PENS, SEA TURTLES DO NOT FACE WITH PEOPLE AND MARINE VEHICLE AND THEY ARE KEPT IN A REDUCED STRESS CONDITIONS.



## WHAT TO DO WHEN AN INJURED OR DEAD TURTLE IS ENCOUNTERED



During on-site controls, it is very important to understand the situation of a turtle that drifts on the sea surface or is stranded to the shore. If the condition of turtle is well examined, the effectiveness of following process will be increased.



When a drifting turtle is encountered on the surface of the sea, it must be taken onboard carefully and transported to a safe place.



Turtle should not be exposed to sun light to prevent rapid body temperature change and should be kept wet to prevent dehydration.



The species of the turtle identified and if possible, measurements should be taken (measurement methodology was described in methodology). Turtle should be checked for tags. If a tag is found, the serial number and the address should be noted.



If turtle is injured, wound area should be identified rapidly and the wound should be cleaned. If this process takes a long time, turtle can be fed.



If a fishing-line is detected, line should not be pulled. If the turtle is found in a remote area with a fishing-line or a hook and the turtle looks healthy, line and hook can be cut after taking measurements and tag controls.



Turtles do not have teeth but they have very strong jaws. Do not put your hand into their mouth.

## WHAT TO DO WHEN AN INJURED OR DEAD TURTLE IS ENCOUNTERED



If the hook is ingested and hooked to esophagus; If the hook is visible, mouth can be open with a hard material and the hook can be cut and removed from the esophagus. Sharp hook should be cut from the feathered edge. A strong pliers can be used to cut the hook. After this operation, turtle should be released when it looks vigorous.



If turtle is not active, it should be kept in a shaded and wet environment. Back of the turtle should be lifted about 20 cm and kept at least 30 minute in this situation. If turtle is not active, it should not be put in the water. If you rescued the turtle from drowning, putting back in the water may cause death. Lifting posterior part of the turtle help removing water from their lungs.



During these process, you can touch turtle's eyelid to check its reflex. If turtle reacts, you have an alive one. Sometimes it may take 24 hours to recover and get reflex. If turtle is not responding and body starts to harden, turtle is dead.



Sometimes turtles can be entangled to trawl nets. If you notice a sea turtle in the trawl net, be careful while depositing the net. If turtle is active, you can release.



In some cases, turtle can be found with small scratches. If turtle is active and wounds are on the body surface, you can release the turtle. If turtle has a head damage, you should keep it. You may apply antibiotic ointment regularly.

## WHAT TO DO WHEN AN INJURED OR DEAD TURTLE IS ENCOUNTERED



If you have to keep turtle for treatment, food and water needs should be met. If turtle starts to lift its head while breathing, it is a good sign. When you hold its flippers, if turtle resist this is again a good sign about its health condition. Another signs of a good health are; when you put turtle on the floor, if it starts to crawl or when you lift in the air if it starts to flap its flipper continuously you can say the turtle is healthy. You can release the turtle as soon as possible if turtle shows these signs.



When you lift the turtle, if the head and the limbs fall and it does not move, it means turtle is injured or, sick or dead. You should check its reflex to understand its state. If turtle is not reacting anything, you can touch its tail (cloacal).



If the turtle is swelled and different animals started to grow on its body, turtle is already dead. You can check for tags, tags have very valuable information for their conservation.



Date, location, species name, genders, ages (juvenile, subadult or adult) and morphological deformation can be seen at first glance of dead turtles should be recorded, and If possible, body size measurements such as curved carapace length and width using a tape measure from the longest parts of the carapace should be taken.



## RECOMMENDATIONS FOR USING TECHNIQUES

WHILE APPLYING THE TECHNIQUES, EXAMINE THE SEA TURTLE CLOSELY, HOLD IT TIGHT BUT GENTLY WITHOUT INJURING OR DAMAGING IT. THE TECHNIQUES CAN BE USED QUICKLY, BUT CAN BE REPEATED TWICE SUCCESSIVELY IF THE FIRST ONE IS SUCCESSFUL. IF THERE IS NO CERTAINTY ABOUT TURTLE'S REACTIONS, WAIT 10 MINUTES BETWEEN THE FIRST AND SECOND TRIAL.





If a turtle is found in the net, lift the posterior



If turtle is active, hold the turtle from the front flippers as it is seen in the figure to stabilize for examining. Do not hold from the tip of the flippers



Turtle shell should be measured with a tape measure as it is seen in the figure



4



If turtle is still, you can touch its eyelid as it is seen in the figure



5



If you don't get a response from the eye, you can touch its tail to get response







## REFERENCES

- Baran, I. and Kasparek, M., 1989. Marine turtles Turkey, status survey 1988 and recommendations for conservation and management. Max Kasparek Verlag, Heidelberg, 123 p.
- Baran, I., Durmuş, S. H., Türközan, O., 1998. Erster nachweis der lederschildekröte, *Dermocheilus coriacea* [Linnaeus, 1766] (Testudines: Dermochelidae) aus Türkischen gewässern. *Herpetofauna*, 20 (1/2): 34–37.
- Baran, I., 1990. Sea turtles in Turkey. *MTP*, 48, 21–22.
- Baran, I., Durmuş, H., Çevik, E., Üçüncü, S. ve Canbolat, R.F., 1992. Türkiye deniz kaplumbağaları stok tesbiti. *Doğa-Tr. J. of Zoology*, 16, 119–139.
- Baran, I., O. Türközan, Ç. Iğaz, Kaska, V. S. Sak, 1996. Research on the marine turtle populations of Dalğan, Fethiye, Patara and Belek Beaches. Final Report, Izmir, 44 p.
- Baran, I., O. Türközan, Ç. Iğaz, S. Sak, N. Taşkın, 1997. Research on the marine turtle populations of Dalğan, Fethiye, Patara and Belek Beaches. Final Report, Izmir, 39 p.
- Başkale, E. ve V. Kaska (2005). Sea turtle nest conservation techniques on Southwestern beaches in Turkey. *Israel Journal of Zoology*, 51, 13–26.
- Başkale, E., Katlımş, V., Rızmaç, M. ve Polat, F. (2012). Fethiye – Göcek Özel Çevre Koruma Bölgesi Tür ve Habitat İzleme Projesi Kapsamında Fethiye Kumsal Alanlarında Deniz Kaplumbağaları (Caretta caretta, Chelonia mydas) Populasyonlarının Araştırılması İzlenmesi ve Korunması. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Başkale, E., Katlımş, V., Rızmaç, M., Yaka, H., Çapar, D., Kaska, V. (2013). Monitoring and Conservation of Sea Turtles (*C. caretta*, *C. mydas*) Populations within the Scope of Fethiye-Göcek Specially Protected Area Monitoring Species and Habitat Project. General Director of the General Directorate of the Protection of Natural Assets, Ministry of Environment and Urbanization, Ankara.
- Başkale, E., Katlımş, V., Rızmaç, M., Sozbielen, D., Polat, F., Lambropoulos, M., Stachowitsch, M. and Kaska, V. (2016). Monitoring and Conservation of Loggerhead Turtle's nests on Fethiye Beaches, Turkey. *Biharean Biologist*, 10(1):20–23.
- Başoğlu, M. (1973). A Preliminary Report on a Specimen of Soft-Shell Turtle from Southwestern Anatolia. *Ege Üniversitesi Fen Fakültesi İlimi Rapor Ser. No.*, 172, 1–11.
- Başoğlu, M., 1973. Sea turtles and the species found along the coasts of neighboring countries. *Türk Biyoloji Dergisi*, 23, 12–21.
- Başoğlu, M. ve Baran, I., 1982. Anadolu sahillerinde toplanan deniz kaplumbağası materyali üzerine kısa bir rapor. *Doğa, Temel Bilimler, Seri R. 6*, 2, 69–71.
- Canbolat, R.F., 1991. Dalğan Kumsalı (Muğla, Türkiye)nda *Caretta caretta* (LINNAEUS 1758) popülasyonu üzerine incelemeler. *Doğa-Tr. J. of Zoology*, 15, 255–274.
- Canbolat, R.F., 1997. Dalğan ve Patara *Caretta caretta* (Linnaeus 1758) Deniz Kaplumbağası Populasyonlarının Biyolojisi. *DÖKTURH TEZ*, HU FBE, Ankara, 454 sayfa.
- Canbolat, R.F., 1999. Köyceğiz-Dalğan ve Patara Özel Çevre Koruma Bölgelerindeki kumsallarda deniz kaplumbağalarının popülasyonlarının araştırılması. *ÖÇKKB Sonuç Raporu*, Ankara, 73 s.
- Canbolat, R.F., 2001c. Köyceğiz-Dalğan Özel Çevre Koruma Bölgesi'nde 2000 ve 2001 Üzere Sezonundaki Deniz Kaplumbağası Populasyonunun Araştırılması. Sonuç Raporu, Hacettepe Üniversitesi-Biyoloji Bölümü, Ankara, 68 s.
- Canbolat, R.F., 2002. Belek ve Köyceğiz-Dalğan Özel Çevre Koruma Bölgelerinde Deniz Kaplumbağaları İzleme Projesi, 2002 Yılı Sonuç Raporu, Özel Çevre Koruma Kurumu Başkanlığı Ankara, 152 s.
- Canbolat, R.F., 2003. Köyceğiz-Dalğan Özel Çevre Koruma Bölgesi'ndeki Deniz Kaplumbağası Populasyonlarının İzlenmesi ve Korunması Projesi. Destekleyen, Çevre ve Orman Bakanlığı-Özel Çevre Koruma Kurumu Başkanlığı, Sürüten Kuruluş: Ekolojik Araştırmalar Derneği, Ankara.
- Canbolat R.F. (2004). A review of sea turtle nesting activity along the Mediterranean coast of Turkey. *Biol Conserv* 116: 81–91.
- Canbolat, R.F., 2006. Köyceğiz-Dalğan ve Fethiye Özel Çevre Koruma Bölgelerinde Deniz Kaplumbağası (Caretta caretta, Chelonia mydas) ve Nil Kaplumbağası (Trionyx triunguis) Populasyonlarının Araştırılması ve Korunması Projesi Hizmet İlimi İş Projesi. Destekleyen, Çevre ve Orman Bakanlığı-Özel Çevre Koruma Kurumu Başkanlığı, Muğla Özel Çevre Koruma Müdürlüğü, Muğla, 148 s.
- Canbolat, R.F., 2007. Köyceğiz-Dalğan ve Fethiye Özel Çevre Koruma Bölgelerinde Deniz Kaplumbağası (Caretta caretta, Chelonia mydas) ve Nil Kaplumbağası (Trionyx triunguis) Populasyonlarının Araştırılması ve Korunması Projesi Hizmet İlimi İş Projesi. Destekleyen, Çevre ve Orman Bakanlığı-Özel Çevre Koruma Kurumu Başkanlığı, Muğla Özel Çevre Koruma Müdürlüğü, Muğla, 148 s.
- Canbolat, R.F., Rızaç, Z., Hoş, R. C., Metin, H., Akbaba, B., 2010. Fethiye-Göcek Özel Çevre Koruma Bölgesinde Deniz Kaplumbağası (Caretta caretta, Chelonia mydas) ve Nil Kaplumbağası (Trionyx triunguis) Populasyonlarının Araştırılması ve Korunması. T.C. Çevre ve Orman Bakanlığı, Özel Çevre Koruma Kurumu Başkanlığı, Ankara.
- Durmuş, H. ve Güçlü, D. (2009). Köyceğiz-Dalğan Özel Çevre Koruma Bölgesi, Dalğan (Iztuz) Kumsal Alanında 2009 Yılı İçin Deniz Kaplumbağaları (Caretta caretta, Chelonia mydas) ve Nil Kaplumbağası (Trionyx triunguis) Populasyonlarının Korunması Ve İzlenmesi Projesi. T.C. Çevre ve Orman Bakanlığı, Özel Çevre Koruma Kurumu Başkanlığı, Ankara.
- Erkakan, F., 1993. Nesting biology of loggerhead turtles *Caretta caretta* L. on Dalğan Beach, Muğla-Turkey. *Biological Conservation*, 66, 1–4.
- Getliş, R. ve Koraç, T., 1982. Türkiye'nin Ege ve Akdeniz kıyılarında yaşayan deniz kaplumbağalarının (*Caretta c. caretta* L. ve *Chelonia m. mydas* L.) popülasyonları ve korunmaları ile ilgili tedbirler üzerine araştırmalar. TÜBİTAK Proje no: UHKG-431, 121 s.
- Geldiği, R., 1983. Deniz kaplumbağalarının (Caretta c. caretta L. ve *Chelonia m. mydas* L.) korunmasında temel bilimler yönünden takip edilecek stratejinin önemi. E. Ü. Fen Fak. Dergisi, Seri B, 1, 328–349.

- Geldiay, R., 1984. Türkiye'nin Ege ve Akdeniz kıyılarında yaşayan deniz kaplumbağalarının (*Caretta c. caretta* L. ve *Chelonia m. mydas* L.) populasyonları ve korunması ile ilgili araştırmalar, Doğa Bilim Dergisi, 82, 1. 66-75.
- Geldiay, R., Koray, T. and Balk, S., 1982. Status of the sea turtle population (*Caretta c. caretta* and *Chelonia m. mydas*) in the Northern Mediterranean Sea, Turkey, p. 424-435, in: K.F. Bjorndal [Ed.], *Biology and Conservation of Sea Turtles*, Washington, DC, 583 p.
- Groombridge, B., 1990. Marine turtles in the Mediterranean; distribution, population status, conservation. Report to the Council of Europe, World Conservation Monitoring Centre, Cambridge, UK, 72 p.
- Hathaway, R.R., 1972. Sea turtles, unanswered questions about sea turtles in Turkey, Balk ve Balkçılık, 20, 1, 1-8.
- Hoofien, J.H. (1972). Reptiles of Israel. Dept. of Zoology, Tel Aviv University, Israel.
- Hoş R.C., Metin H., Bulut Ş., Candan D., 2014. Fethiye-Göcek Özel Çevre Koruma Bölgesi Tür ve Habitat İzleme Projesi Kapsamında Fethiye Kumsal Alanında Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Araştırılması, İzlenimi ve Korunması Projesi Sonuç Raporu, T.C Çevre ve Şehircilik Bakanlığı, Tabiat Varlıklarını Koruma Genel Müdürlüğü, Ankara, 2014, 114 s.
- Ilgaz, C., Baran, L., 2001. Reproduction biology of the marine turtle populations in Northern Karpaz (Northern Cyprus) and Dalğın (Turkey). *Zoology in the Middle East* 24, 35-44.
- Ilgaz, C., Türkozan, D., Özdemir, R., Kaska, Y., Stachowiutz, M., 2007. Population Decline of Loggerhead Turtles: Two Potential Scenarios for Fethiye Beach, Turkey. *Biodiversity and Conservation* 16(4): 1027-1037.
- Kaska, Y., 1993. Kızlot ve Patara *Caretta caretta* populasyonunun araştırılması. Master Tezi, D.E.Ü. Fen Bilimleri Enstitüsü, İzmir, 28 s.
- Kaska, Y., Dounie, J.R., Tippett, R. ve Furness, R. (1998). Natural Temperature Regimes for Loggerhead and Green Turtle Nests in the Eastern Mediterranean, *Can. J. Zool.*, 76, 723-729.
- Kaska, Y., İlbaran, Ç., Ilgaz, O., Türkozan, M., Öz ve R. Erdogan, "An estimation of the total nesting activity of sea turtles in Turkey", Pp. 204-205 In: Coyne P.S., and Clark, R.D. (Compilers), *Proceedings of the Twenty-First Annual Sea Turtle Biology and Conservation*. NOAA Technical Memorandum NMFS-SEFCS-528, 368 pp. (İlami, 2005).
- Kaska, Y., Ilgaz, C., Özdemir, R., Başkale, E., Türkozan, D., Baran, L. and Stachowiutz, M. (2006). Sex ratio estimations of loggerhead sea turtle hatchlings by histological examination and nest temperatures at Fethiye beach, Turkey. *Naturwissenschaften*, 93: 338-343.
- Kaska, Y., Sözbilen, D. ve Sarı, F. (2008). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2008 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Orman Bakanlığı, Özel Çevre Koruma Kurumu Başkanlığı, Ankara. 978-9944-0847-1-0.
- Kaska, Y., Dişen, S., Başkale, E., Sarı, F. ve Ulubeli, R.S. (2010). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2010 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Orman Bakanlığı, Özel Çevre Koruma Kurumu Başkanlığı, Ankara.
- Kaska, Y., Başkale, E., Fak, Ç. (2011). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2011 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Başkale, E., ve Kara, I. (2011). Fethiye - Göcek Özel Çevre Koruma Bölgesi Tür ve Habitat İzleme Projesi Kapsamında Fethiye Kumsal Alanlarında Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) Populasyonlarının Araştırılması İzlenimi ve Korunması. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Dişen, S., Fak, Ç., Yaka, H. (2012). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2012 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Dişen, S., Sözbilen, D., Sezgin, Ç. (2013). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2013 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Sözbilen, D., Sezgin, Ç., Candan, R.Y. (2014). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2014 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Başkale, E., Sözbilen, D., Sezgin, Ç., Candan, R.Y. (2015). Köyceğiz-Dalğın Özel Çevre Koruma Bölgesi, Dalğın İztuzu Kumsal Alanında 2015 Yılı İçin Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Korunması ve İzlenimi Projesi. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Başkale, E., Katlımş, Y., Sözbilen, D., Rızmaz, M. ve Candan, R.Y. 2016. Köyceğiz-Dalğın Kumsal Alanında Deniz Kaplumbağası (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Araştırılması İzlenimi ve Korunması Projesi. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Kaska, Y., Başkale, E., Katlımş, Y., Sözbilen, D., Rızmaz, M. ve Üzulmaz, Y. 2016. Fethiye - Göcek Özel Çevre Koruma Bölgesi Tür ve Habitat İzleme Projesi Kapsamında Fethiye Kumsal Alanlarında Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) Populasyonlarının Araştırılması İzlenimi ve Korunması. T.C. Çevre ve Şehircilik Bakanlığı, Ankara.
- Lutz, P.L. ve Musick, J.R. (1997). *The Biology of Sea Turtles*, CRC Press, New York, 432 pp.
- Kaska, Y., Başkale, E., Urhan, R., Katlımş, Y., Gidiş, M., Sarı, F., Sözbilen, D., Canbolat, R.F., Vilmaz, F., Barlas, M., Özdemir, N., ve Özkul (2010). Natural and anthropogenic factors affecting the nest-site selection of Loggerhead Turtles, *Caretta caretta*, on Dalaman-Sarıgerme beach in South-west Turkey. *Zoology in the Middle East*, 50:47-58.
- Ozner, S. 1991. Çarpık yapılaşma ve aşırı kum alımları Türkiye'nin Akdeniz kıyı kuşağında kuyi dengesi ve deniz kaplumbağaları üzerindeki olumsuz etkileri, *Jeomorfoloji Dergisi*, 19, 151-158.
- Ripple, J. 1996. *Sea turtles*. Stillwater, USA: Voyageur Press, Inc.
- Sönmez, B., Sumnu, D., Valçın Özdelek, Ş., Gonenler, Ö.R., Açıkgöz, U., Ergün, Y., Kaska, Y. 2008. A stranded leatherback sea turtle (*Dermochelys coriacea* Vandelli, 1761) in the Northeastern Mediterranean. *Hatay Turkey Marine Turtle Newsletter*, 19(1), 12-13.
- TRKUR, (2008). Fethiye-Göcek Özel Çevre Koruma Bölgelerinde Deniz Kaplumbağası (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının İzlenimi ve Korunması Projesi Hizmet Filmi İş Projesi. Destekleyen, Çevre ve Orman Bakanlığı-Özel Çevre Koruma Kurumu Başkanlığı, Ankara, 55 s.
- TRKUR, (2009). Fethiye-Göcek Özel Çevre Koruma Bölgelerinde Deniz Kaplumbağası (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının İzlenimi ve Korunması Projesi Hizmet Filmi İş Projesi. Destekleyen, Çevre ve Orman Bakanlığı-Özel Çevre Koruma Kurumu Başkanlığı, Ankara, 76 s.
- Taşkavak, E., Boulon, R.H. Attard, M.K., 1998. An Unusual Stranding of a Leatherback Turtle in Turkey. *Marine Turtle Newsletter* (80):13.
- Türkozan, D., 2006. Preliminary results for the population trends of two index beaches, Fethiye and Dalğın. In: Frick (Ed), Panagopolou R, Rees AF, Williams M (eds) *Twenty sixth Annual Symposium on Sea Turtle Biology and Conservation*, Athens, Greece, 376 p. 330.
- Türkozan, D., Vilmaz, C., 2008. Loggerhead turtles, *Caretta caretta*, at Dalğın Beach, Turkey: nesting activity (2004-2005) and 19-year abundance trend (1987-2005). *Chelonian Conserv* 8(7): 178-187.
- Türkozan, D. and Baran, L., 1996. Research on the loggerhead turtle, *Caretta caretta*, of Fethiye Beach, Tr. *J. of Zoology*, 20, 183-188.
- Türkozan, D. (2000). Reproductive Ecology of the Loggerhead Turtle, *Caretta caretta*, on Fethiye and Kızlot Beaches, Turkey. *Chelonian Cons. and Biol.* 6:86-92.
- Türkozan, D., Ilgaz, C., Taşkavak, E., Özdemir, R., 2003. Hatch rates of loggerhead turtles and physical characteristics of the beach at Fethiye. *Journal of the Marine Biological Association, U.K.* 83: 231-232
- Türkozan ve Kaska (2010). Turkey: In: Sea Turtles in the Mediterranean: Distribution, Threats and conservation priorities (Casale and Margaritoulis [Eds]), Gland, Switzerland, IUCN, p 257-294.
- Yerli, S.M., F. Demirayak, 1996. Türkiye'de Deniz Kaplumbağaları ve Üreme Kumsalları Üzerine Bir Değerlendirme '95. DHKD-Kıyı Yönetimi Bölümü 96/4, İstanbul, 129 s.
- Yerli, S.M., Mangıt, F., Korkmaz, M., Mantarlı, S., Sü, U., İçsen, B. ve Güreşoğlu, C., 2015. Fethiye - Göcek Özel Çevre Koruma Bölgesi Tür ve Habitat İzleme Projesi Kapsamında Fethiye Kumsal Alanında Deniz Kaplumbağaları (*Caretta caretta*, *Chelonia mydas*) ve Fıllı Kaplumbağası (*Trionyx triunguis*) Populasyonlarının Araştırılması, İzlenimi ve Korunması Projesi Sonuç Raporu, T.C. Çevre ve Şehircilik Bakanlığı, Tabiat Varlıklarını Koruma Genel Müdürlüğü, Ankara, 2015.
- Wltherington, B. E. and Bjorndal, K. R. 1991. Influences of Uvaelength and Intensity on Hatchling Sea Turtle Phototaxis: Implications for Sea- Finding Behavior. *Copeia* 1991: 1060- 1069.



# Caretta caretta



Please Inform Us of Any Injured Sea Turtles...



Sea Turtle Research Rescue and Rehabilitation Centre

Phone / Fax: +90 252 289 00 77 [caretta@pau.edu.tr](mailto:caretta@pau.edu.tr) [www.dekamer.org.tr](http://www.dekamer.org.tr)

---

IF YOU SEE AN INJURED SEA TURTLE PLEASE CALL CLOSEST COAST GUARD OFFICE (112), GENDARMIE (112) OR SEA TURTLE REHABILITATION CENTER -DEKAMER (+90 258 289 0077)  
THANK YOU FOR YOUR COOPERATION

---

**MONITORING AND CONSERVATION STUDIES OF SEA TURTLES (CARETTA CARETTA) DURING THE 2016 NESTING SEASON ON MUĞLA SEA TURTLE NESTING BEACHES**

Kaska, Y., Başkale, E., Katılmış, Y., Sözbilen, D., Azmaz, M. (2016). Monitoring and Conservation Studies of Sea Turtles (*Caretta caretta*) During the 2016 Nesting Season on Muğla Sea Turtle Nesting Beaches . 52 pp. Denizi.



9 786058 1292307