

The Exact Solutions of Conformable Fractional Partial Differential Equations Using New Sub Equation Method

Ali Kurt^{1*}, Orkun Tasbozan² and Hulya Durur³

¹Department of Mathematics, Faculty of Science and Arts, Pamukkale University, Denizli, Turkey

²Department of Mathematics, Faculty of Science and Arts, Hatay Mustafa Kemal University, Hatay, Turkey

³Department of Computer Engineering, Faculty of Engineering, Ardahan University, Ardahan, Turkey

*Corresponding author E-mail: pau.dr.alikurt@gmail.com, akurt@pau.edu.tr

Article Info

Keywords: Conformable fractional derivative, Exact solutions, The new sub-equation method

2010 AMS: 35R11, 35A20, 35C05

Received: 10 May 2019

Accepted: 26 October 2019

Available online: 20 December 2019

Abstract

In this article, authors employed the new sub equation method to attain new traveling wave solutions of conformable time fractional partial differential equations. Conformable fractional derivative is a well behaved, applicable and understandable definition of arbitrary order derivation. Also this derivative obeys the basic properties that Newtonian concept satisfies. In this study authors obtained the exact solution for KDV equation where the fractional derivative is in conformable sense. New solutions are obtained in terms of the generalized version of the trigonometric functions.

1. Introduction

Fractional differential equations (FDEs) are generalized form of the integer order differential equations. In the last decades, researchers have worked hard for obtaining analytic solutions of nonlinear FDEs. Nonlinear FDEs are often used to describe many problems arising in many fields such as physics, chemistry, engineering, heat transfer, applied mathematics, control theory et al. [1]-[4]. So, many authors presented very strong methods to solve FDEs. For instance Kurt et. al. [5] studied the solutions of time fractional Whitham-Broer-Kaup Equation by using homotopy analysis method where the fractional terms are described in Caputo sense. Tasbozan et. al. [6] employed the finite element method for attaining the approximate solutions of diffusion equation where the derivatives are in Riemann-Liouville sense. Celik et. al. [7] utilized Crank-Nicolson scheme to get the the numerical solutions of fractional diffusion equation. As it is seen from the given references, all the obtained results are numerical solutions for the considered nonlinear equations. Because, the analytical methods can not be applied to the nonlinear equations which involves Caputo, Riemann-Liouville and Riesz fractional derivative definitions. On the contrary, conformable fractional derivative definition gives us chance to get the exact solutions of nonlinear FDEs by using new wave transformation [8] and the chain rule [9]. For example Eslami and Rezazadeh [10] used the first integral method to obtain analytic solutions of time fractional Wu-Zhang system. Aminikhah et. al. [11] obtained analytic solutions of fractional regularized long-wave equations using sub-equation method. Osman et al. [12] employed the unified method to get the analytic solutions of conformable time fractional Schrödinger equation with perturbation terms. For further details please see the references [13]-[34]. In this paper, we handle the Korteweg-de Vries equation with a source that provides a sixth order differential equation.

$$D_x^6 u + 20D_x u D_x^4 u + 40D_x^2 u D_x^3 u + 120D_x u^2 D_x^2 u + D_x^3 D_t^\mu u + 8D_x u D_x D_t^\mu u + 4D_t^\mu u D_x^2 u = 0. \quad (1.1)$$

2. Conformable fractional calculus

R. Khalil et. al. [32] presented the definition of conformable fractional derivative as follows.

Definition 2.1. μ^{th} order "conformable fractional derivative" of function g which is defined as $g : [0, \infty) \rightarrow \mathbb{R}$ can be dedicated as

$$T_\mu(g)(t) = \lim_{\varepsilon \rightarrow 0} \frac{g(t + \varepsilon t^{1-\mu}) - g(t)}{\varepsilon}$$

for all $t > 0$, $\alpha \in (0, 1)$. Assuming that g is μ -differentiable over some $(0, a)$ where $a > 0$ and $\lim_{t \rightarrow 0^+} g^{(\mu)}(t)$ exists, then $g^{(\mu)}(0) = \lim_{t \rightarrow 0^+} g^{(\mu)}(t)$.

The other fractional derivative definitions such as Caputo, Riemann-Liouville, Grünwald-Letnikov and etc. do not satisfy basic principles which are provided by Newtonian type derivative. For instance

1. Assume that λ is a constant and $\alpha \in \mathbb{R}$. Then $D_a^\mu(\lambda) \neq 0$ for Riemann-Liouville derivative.
2. The Riemann-Liouville and Caputo derivatives do not provide the derivative of the product of two functions.
3. $D_a^\mu(fg) \neq fD_a^\mu(g) + gD_a^\mu(f)$.
4. The Riemann-Liouville and Caputo derivatives do not provide the derivative of the quotient of two functions
5. $D_a^\mu\left(\frac{f}{g}\right) \neq \frac{gD_a^\mu(f) - fD_a^\mu(g)}{g^2}$.

This new definition satisfies the properties which are given in the following theorem.

Theorem 2.2. Let $\mu \in (0, 1)$ and f, g be μ -differentiable at point $t > 0$. Then

1. $T_\mu(af + bg) = aT_\mu(f) + bT_\mu(g)$, for all $a, b \in \mathbb{R}$
2. $T_\mu(t^p) = pt^{p-\mu}$ for all $p \in \mathbb{R}$.
3. $T_\mu(\lambda) = 0$ for all constant function $f(t) = \lambda$.
4. $T_\mu(fg) = fT_\mu(g) + gT_\mu(f)$.
5. $T_\mu\left(\frac{f}{g}\right) = \frac{gT_\mu(f) - fT_\mu(g)}{g^2}$.
6. If f is differentiable, then $T_\mu(f)(t) = t^{1-\mu} \frac{df}{dt}$.

3. The new sub-equation method

Consider that the general form of nonlinear fractional partial differential equation can be expressed as

$$H\left(u, \frac{\partial^\mu u}{\partial t^\mu}, \frac{\partial u}{\partial x}, u \frac{\partial u}{\partial x}, u^2 \frac{\partial u}{\partial x}, \frac{\partial^2 u}{\partial x^2}, \dots\right) = 0. \quad (3.1)$$

Using the wave transform $\xi = kx + w \frac{t^\mu}{\mu}$ where k and w are constants and chain rule [9] in Eq. (3.1), the independent variables and can be changed into single variable. So Eq. (3.1) can be rewritten as

$$P(u, u'(\xi), u''(\xi), \dots). \quad (3.2)$$

Consider that $u(\xi)$ can be written as a polynomial in $Q(\xi)$

$$u(\xi) = \sum_{j=0}^n a_j Q^j(\xi), \quad (3.3)$$

where a_j ($0 \leq j \leq n$) are constant coefficients to be determined after and $Q(\xi)$ provides first order linear ODE of the form

$$Q'(\xi) = Ln(A) \left(\alpha + \beta Q(\xi) + \sigma Q^2(\xi) \right), \quad A \neq 0, 1, \quad (3.4)$$

where α, β, σ are constants. Moreover, Eq. has the following traveling wave solutions.

Family 1. If $\beta^2 - 4\alpha\sigma < 0$ and $\sigma \neq 0$, then we have

$$\begin{aligned} Q_1(\xi) &= -\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \tan_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2} \xi \right), \\ Q_2(\xi) &= -\frac{\beta}{2\sigma} - \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \cot_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2} \xi \right), \\ Q_3(\xi) &= -\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \left(\tan_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \pm \sqrt{pq} \sec_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \right), \\ Q_4(\xi) &= -\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \left(-\cot_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \pm \sqrt{pq} \csc_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \right), \\ Q_5(\xi) &= -\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{4\sigma} \left(\tan_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{4} \xi \right) - \cot_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{4} \xi \right) \right). \end{aligned}$$

Family 2. Suppose that $\beta^2 - 4\alpha\sigma > 0$ and $\sigma \neq 0$,

$$\begin{aligned} Q_6(\xi) &= -\frac{\beta}{2\sigma} - \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \tanh_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2} \xi \right), \\ Q_7(\xi) &= -\frac{\beta}{2\sigma} - \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \coth_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2} \xi \right), \\ Q_8(\xi) &= -\frac{\beta}{2\sigma} + \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \left(-\tanh_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \pm i\sqrt{pq} \operatorname{sech}_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \right), \\ Q_9(\xi) &= -\frac{\beta}{2\sigma} + \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \left(-\coth_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \pm \sqrt{pq} \operatorname{csch}_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \right), \\ Q_{10}(\xi) &= -\frac{\beta}{2\sigma} - \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{4\sigma} \left(\tanh_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{4} \xi \right) + \coth_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{4} \xi \right) \right). \end{aligned}$$

Family 3. Consider that $\alpha\sigma > 0$ and $\beta = 0$,

$$\begin{aligned} Q_{11}(\xi) &= \sqrt{\frac{\alpha}{\sigma}} \tan_A (\sqrt{\alpha\sigma} \xi), \\ Q_{12}(\xi) &= -\sqrt{\frac{\alpha}{\sigma}} \cot_A (\sqrt{\alpha\sigma} \xi), \\ Q_{13}(\xi) &= \sqrt{\frac{\alpha}{\sigma}} \left(\tan_A (2\sqrt{\alpha\sigma} \xi) \pm \sqrt{pq} \sec_A (2\sqrt{\alpha\sigma} \xi) \right), \\ Q_{14}(\xi) &= \sqrt{\frac{\alpha}{\sigma}} \left(-\cot_A (2\sqrt{\alpha\sigma} \xi) \pm \sqrt{pq} \csc_A (2\sqrt{\alpha\sigma} \xi) \right), \\ Q_{15}(\xi) &= \frac{1}{2} \sqrt{\frac{\alpha}{\sigma}} \left(\tan_A \left(\frac{\sqrt{\alpha\sigma}}{2} \xi \right) - \cot_A \left(\frac{\sqrt{\alpha\sigma}}{2} \xi \right) \right). \end{aligned}$$

Family 4. Regard that $\alpha\sigma < 0$ and $\beta = 0$,

$$\begin{aligned} Q_{16}(\xi) &= -\sqrt{-\frac{\alpha}{\sigma}} \tanh_A (\sqrt{-\alpha\sigma} \xi), \\ Q_{17}(\xi) &= -\sqrt{-\frac{\alpha}{\sigma}} \coth_A (\sqrt{-\alpha\sigma} \xi), \\ Q_{18}(\xi) &= \sqrt{-\frac{\alpha}{\sigma}} \left(-\tanh_A (2\sqrt{-\alpha\sigma} \xi) \pm i\sqrt{pq} \operatorname{sech}_A (2\sqrt{-\alpha\sigma} \xi) \right), \\ Q_{19}(\xi) &= \sqrt{-\frac{\alpha}{\sigma}} \left(-\coth_A (2\sqrt{-\alpha\sigma} \xi) \pm \sqrt{pq} \operatorname{csch}_A (2\sqrt{-\alpha\sigma} \xi) \right), \\ Q_{20}(\xi) &= -\frac{1}{2} \sqrt{-\frac{\alpha}{\sigma}} \left(\tanh_A \left(\frac{\sqrt{-\alpha\sigma}}{2} \xi \right) + \coth_A \left(\frac{\sqrt{-\alpha\sigma}}{2} \xi \right) \right). \end{aligned}$$

Family 5. When $\beta = 0$ and $\sigma = \alpha$,

$$\begin{aligned} Q_{21}(\xi) &= \tan_A (\alpha\xi), \\ Q_{22}(\xi) &= -\cot_A (\alpha\xi), \\ Q_{23}(\xi) &= \tan_A (2\alpha\xi) \pm \sqrt{pq} \sec_A (2\alpha\xi), \\ Q_{24}(\xi) &= -\cot_A (2\alpha\xi) \pm \sqrt{pq} \csc_A (2\alpha\xi), \\ Q_{25}(\xi) &= \frac{1}{2} \left(\tan_A \left(\frac{\alpha}{2} \xi \right) - \cot_A \left(\frac{\alpha}{2} \xi \right) \right). \end{aligned}$$

Family 6. If $\beta = 0$ and $\sigma = -\alpha$, chosen

$$\begin{aligned} Q_{26}(\xi) &= -\tanh_A (\alpha\xi), \\ Q_{27}(\xi) &= -\coth_A (\alpha\xi), \\ Q_{28}(\xi) &= -\tanh_A (2\alpha\xi) \pm i\sqrt{pq} \operatorname{sech}_A (2\alpha\xi), \\ Q_{29}(\xi) &= -\coth_A (2\alpha\xi) \pm \sqrt{pq} \operatorname{csch}_A (2\alpha\xi), \\ Q_{30}(\xi) &= -\frac{1}{2} \left(\tanh_A \left(\frac{\alpha}{2} \xi \right) + \coth_A \left(\frac{\alpha}{2} \xi \right) \right). \end{aligned}$$

Family 7. While $\beta^2 = 4\alpha\sigma$,

$$Q_{31}(\xi) = \frac{-2\alpha(\beta\xi \operatorname{Ln}(A) + 2)}{\beta^2 \xi \operatorname{Ln}(A)}.$$

Family 8. When $\beta = k$, $\alpha = mk (m \neq 0)$ and $\sigma = 0$,

$$Q_{32}(\xi) = A^{k\xi} - m.$$

Family 9. When $\beta = \sigma = 0$,

$$Q_{33}(\xi) = \alpha \xi \operatorname{Ln}(A).$$

Family 10. When $\beta = \alpha = 0$,

$$Q_{34}(\xi) = \frac{-1}{\sigma \xi \operatorname{Ln}(A)}.$$

Family 11. When $\alpha = 0$ and $\beta \neq 0$,

$$\begin{aligned} Q_{35}(\xi) &= -\frac{p\beta}{\sigma(\cosh_A(\beta\xi) - \sinh_A(\beta\xi) + p)}, \\ Q_{36}(\xi) &= -\frac{q\beta}{\sigma(\cosh_A(\beta\xi) - \sinh_A(\beta\xi) + q)}, \\ Q_{37}(\xi) &= -\frac{\beta(\sinh_A(\beta\xi) + \cosh_A(\beta\xi))}{\sigma(\sinh_A(\beta\xi) + \cosh_A(\beta\xi) + q)}, \end{aligned}$$

Family 12. When $\beta = k$, $\sigma = mk (m \neq 0)$, $p = q$ and $\alpha = 0$,

$$Q_{38}(\xi) = \frac{pA^{k\xi}}{p - mqA^{k\xi}}.$$

Remark 3.1. The generalized version of the trigonometric functions and the generalized types of the hypergeometric functions are declared as [33]

$$\begin{aligned} \sinh_A(\xi) &= \frac{pA^\xi - qA^{-\xi}}{2}, & \cosh_A(\xi) &= \frac{pA^\xi + qA^{-\xi}}{2}, \\ \tanh_A(\xi) &= \frac{pA^\xi - qA^{-\xi}}{pA^\xi + qA^{-\xi}}, & \coth_A(\xi) &= \frac{pA^\xi + qA^{-\xi}}{pA^\xi - qA^{-\xi}}, \\ \operatorname{sech}_A(\xi) &= \frac{2}{pA^\xi + qA^{-\xi}}, & \operatorname{csch}_A(\xi) &= \frac{2}{pA^\xi - qA^{-\xi}}, \\ \sin_A(\xi) &= \frac{pA^{i\xi} - qA^{-i\xi}}{2i}, & \cos_A(\xi) &= \frac{pA^{i\xi} + qA^{-i\xi}}{2}, \\ \tan_A(\xi) &= -i \frac{pA^{i\xi} - qA^{-i\xi}}{pA^{i\xi} + qA^{-i\xi}}, & \cot_A(\xi) &= i \frac{pA^{i\xi} + qA^{-i\xi}}{pA^{i\xi} - qA^{-i\xi}}, \\ \sec_A(\xi) &= \frac{2}{pA^{i\xi} + qA^{-i\xi}}, & \operatorname{csc}_A(\xi) &= \frac{2i}{pA^{i\xi} - qA^{-i\xi}}, \end{aligned}$$

where $p, q > 0$ are constants and ξ is an independent variable. In addition, by considering the balance between the highest order derivative linear term and nonlinear terms appearing in ODE (3.2), the positive integer n can be defined. Replacing Eq. (3.3) into ODE (3.2), using Eq. (3.4), and equalizing the coefficients of all the powers of $Q(\xi)$ to zero, we will obtain an equation system in terms of k, w and $a_j (0 \leq j \leq n)$. From this obtained system the values for k, w and a_j can be found with the aid of a computer software. Replacing the obtained values of k, w and a_j into Eq.(3.3), we may acquire all possible solutions of Eq. (3.1).

4. Analytic results for time fractional KdV6 equation with conformable derivative

Using the wave transformation and applying chain rule [9]

$$u(x, t) = u(\xi), \quad \xi = kx + w \frac{t^\mu}{\mu}. \tag{4.1}$$

Eq. (1.1) is transferred to

$$k^6 u^{(vi)}(\xi) + k^3 w u^{iv}(\xi) + 6k^2 w (u'(\xi))^2 + 20k^5 u^{iv}(\xi) u'(\xi) + 40k^5 u''(\xi) u'''(\xi) + 12k^2 w u'(\xi) u''(\xi) = 0$$

where the prime symbolizes the known derivative of function $u(\xi)$ with respect to ξ . Integrating the above equation once and making some algebraic calculations led to

$$k^6 u^{(v)}(\xi) + k^3 w u'''(\xi) + 3k^2 w (u')^2 + 5k^5 u''' u + 20k^5 (u'')^2 + 12k^2 w u u' = 0. \tag{4.2}$$

Considering the homogeneous balance between $u^2 u'$ and $u^{(5)}$ in Eq. (4.2) we obtain $n + 5 = 3(n + 1)$; then $n = 1$; so we can write Eq. (3.3) as

$$u(\xi) = a_0 + a_1 Q(\xi). \tag{4.3}$$

Subrogating Eq. (4.3) with (3.4) into Eq. (4.2) and gathering all the same power of $Q(\xi)$ together, the left hand side of Eq. (4.2) turns into a polynomial of $Q(\xi)$. Equalizing the each coefficient of the same power of $Q(\xi)$ to zero led to an equation system. Solving the obtained system due to unknowns variables a_0 , a_1 and w , the solutions can be concluded as

$$w = \frac{a_1^3 (\beta^2 - 4\alpha\sigma)}{\sigma^3 Ln(A)}, \quad k = -\frac{a_1}{\sigma Ln(A)}. \tag{4.4}$$

Putting the solution set (4.4) with (4.1) into (4.3) and solutions of Eq. (1.1), can be expressed as

Case 1. If $\beta^2 - 4\alpha\sigma < 0$ and $\sigma \neq 0$, then we have

$$\begin{aligned} u_1(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \tan_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2} \xi \right) \right), \\ u_2(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} - \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \cot_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2} \xi \right) \right), \\ u_3(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \left(\tan_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \pm \sqrt{pq} \sec_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \right) \right), \\ u_4(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{2\sigma} \left(-\cot_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \pm \sqrt{pq} \csc_A \left(\sqrt{-(\beta^2 - 4\alpha\sigma)} \xi \right) \right) \right), \\ u_5(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} + \frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{4\sigma} \left(\tan_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{4} \xi \right) - \cot_A \left(\frac{\sqrt{-(\beta^2 - 4\alpha\sigma)}}{4} \xi \right) \right) \right) \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma Ln(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu Ln(A)}t^\mu$.

Case 2. Suppose that $\beta^2 - 4\alpha\sigma > 0$ and $\sigma \neq 0$,

$$\begin{aligned} u_6(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} - \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \tanh_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2} \xi \right) \right), \\ u_7(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} - \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \coth_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2} \xi \right) \right), \\ u_8(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} + \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \left(-\tanh_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \pm i\sqrt{pq} \operatorname{sech}_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \right) \right), \\ u_9(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} + \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{2\sigma} \left(-\coth_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \pm \sqrt{pq} \operatorname{sch}_A \left(\sqrt{\beta^2 - 4\alpha\sigma} \xi \right) \right) \right), \\ u_{10}(\xi) &= a_0 + a_1 \left(-\frac{\beta}{2\sigma} - \frac{\sqrt{\beta^2 - 4\alpha\sigma}}{4\sigma} \left(\tanh_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{4} \xi \right) + \coth_A \left(\frac{\sqrt{\beta^2 - 4\alpha\sigma}}{4} \xi \right) \right) \right) \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma Ln(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu Ln(A)}t^\mu$.

Case 3. Consider that $\alpha\sigma > 0$ and $\beta = 0$,

$$\begin{aligned} u_{11}(\xi) &= a_0 + a_1 \left(\sqrt{\frac{\alpha}{\sigma}} \tan_A \left(\sqrt{\alpha\sigma} \xi \right) \right), \\ u_{12}(\xi) &= a_0 - a_1 \left(\sqrt{\frac{\alpha}{\sigma}} \cot_A \left(\sqrt{\alpha\sigma} \xi \right) \right), \\ u_{13}(\xi) &= a_0 + a_1 \left(\sqrt{\frac{\alpha}{\sigma}} \left(\tan_A \left(2\sqrt{\alpha\sigma} \xi \right) \pm \sqrt{pq} \sec_A \left(2\sqrt{\alpha\sigma} \xi \right) \right) \right), \\ u_{14}(\xi) &= a_0 + a_1 \left(\sqrt{\frac{\alpha}{\sigma}} \left(-\cot_A \left(2\sqrt{\alpha\sigma} \xi \right) \pm \sqrt{pq} \csc_A \left(2\sqrt{\alpha\sigma} \xi \right) \right) \right), \\ u_{15}(\xi) &= a_0 + a_1 \left(\frac{1}{2} \sqrt{\frac{\alpha}{\sigma}} \left(\tan_A \left(\frac{\sqrt{\alpha\sigma}}{2} \xi \right) - \cot_A \left(\frac{\sqrt{\alpha\sigma}}{2} \xi \right) \right) \right) \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma L_n(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu L_n(A)}t^\mu$.

Case 4. Regard that $\alpha\sigma < 0$ and $\beta = 0$,

$$\begin{aligned} u_{16}(\xi) &= a_0 - a_1 \left(\sqrt{-\frac{\alpha}{\sigma}} \tanh_A(\sqrt{-\alpha\sigma\xi}) \right), \\ u_{17}(\xi) &= a_0 - a_1 \left(\sqrt{-\frac{\alpha}{\sigma}} \coth_A(\sqrt{-\alpha\sigma\xi}) \right), \\ u_{18}(\xi) &= a_0 + a_1 \left(\sqrt{-\frac{\alpha}{\sigma}} (-\tanh_A(2\sqrt{-\alpha\sigma\xi}) \pm i\sqrt{pq} \operatorname{sech}_A(2\sqrt{-\alpha\sigma\xi})) \right), \\ u_{19}(\xi) &= a_0 + a_1 \left(\sqrt{-\frac{\alpha}{\sigma}} (-\coth_A(2\sqrt{-\alpha\sigma\xi}) \pm \sqrt{pq} \operatorname{csch}_A(2\sqrt{-\alpha\sigma\xi})) \right), \\ u_{20}(\xi) &= a_0 - a_1 \left(\frac{1}{2} \sqrt{-\frac{\alpha}{\sigma}} \left(\tanh_A\left(\frac{\sqrt{-\alpha\sigma}}{2}\xi\right) + \coth_A\left(\frac{\sqrt{-\alpha\sigma}}{2}\xi\right) \right) \right) \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma L_n(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu L_n(A)}t^\mu$.

Case 5. When $\beta = 0$ and $\sigma = \alpha$,

$$u_{21}(\xi) = a_0 + a_1 \tan_A(\alpha\xi),$$

$$\begin{aligned} u_{22}(\xi) &= a_0 - a_1 \cot_A(\alpha\xi), \\ u_{23}(\xi) &= a_0 + a_1 (\tan_A(2\alpha\xi) \pm \sqrt{pq} \operatorname{sec}_A(2\alpha\xi)), \\ u_{24}(\xi) &= a_0 + a_1 (-\cot_A(2\alpha\xi) \pm \sqrt{pq} \operatorname{csc}_A(2\alpha\xi)), \\ u_{25}(\xi) &= a_0 + a_1 \left(\frac{1}{2} \left(\tan_A\left(\frac{\alpha}{2}\xi\right) - \cot_A\left(\frac{\alpha}{2}\xi\right) \right) \right) \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma L_n(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu L_n(A)}t^\mu$.

Case 6. If $\beta = 0$ and $\sigma = -\alpha$, chosen

$$\begin{aligned} u_{26}(\xi) &= a_0 - a_1 \tanh_A(\alpha\xi), \\ u_{27}(\xi) &= a_0 - a_1 \coth_A(\alpha\xi), \\ u_{28}(\xi) &= a_0 + a_1 (-\tanh_A(2\alpha\xi) \pm i\sqrt{pq} \operatorname{sech}_A(2\alpha\xi)), \\ u_{29}(\xi) &= a_0 + a_1 (-\coth_A(2\alpha\xi) \pm \sqrt{pq} \operatorname{csch}_A(2\alpha\xi)), \\ u_{30}(\xi) &= a_0 - \frac{a_1}{2} \left(\tanh_A\left(\frac{\alpha}{2}\xi\right) + \coth_A\left(\frac{\alpha}{2}\xi\right) \right) \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma L_n(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu L_n(A)}t^\mu$.

Case 11. When $\alpha = 0$ and $\beta \neq 0$,

$$\begin{aligned} u_{31}(\xi) &= a_0 - \frac{pa_1\beta}{\sigma (\cosh_A(\beta\xi) - \sinh_A(\beta\xi) + p)}, \\ u_{32}(\xi) &= a_0 - \frac{qa_1\beta}{\sigma (\cosh_A(\beta\xi) - \sinh_A(\beta\xi) + q)}, \\ u_{33}(\xi) &= a_0 - \frac{a_1\beta (\sinh_A(\beta\xi) + \cosh_A(\beta\xi))}{\sigma (\sinh_A(\beta\xi) + \cosh_A(\beta\xi) + q)}, \end{aligned}$$

where $\xi = -\frac{a_1}{\sigma L_n(A)}x + \frac{a_1^3(\beta^2 - 4\alpha\sigma)^2}{\sigma^3 \mu L_n(A)}t^\mu$.

Case 12. When $\beta = k$, $\sigma = mk (m \neq 0)$, $p = q$ and $\alpha = 0$,

$$u_{34}(\xi) = a_0 + \frac{pa_1 A^{k\xi}}{p - mq A^{k\xi}}.$$

5. Conclusion

In this manuscript the new sub-equation method successfully applied to time fractional KdV6 equation. Analytic solutions of the nonlinear KdV6 equation are successfully obtained. Also wave transform and chain rule are used, so the nonlinear conformable FDE changes into differential equation with integer order derivative. As it can be from the obtained results new sub-equation method is a reliable, efficient and applicable tool for obtaining the exact solutions of fractional partial differential equations in conformable sense.

Acknowledgement

All authors contributed equally to this work.

References

- [1] K. Oldham, J. Spanier, *The Fractional Calculus, Theory and Applications of Differentiation and Integration of Arbitrary Order*, Academic Press, 1974.
- [2] K.S. Miller, B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, A Wiley-Interscience Publication, 1993.
- [3] I. Podlubny, *Fractional Differential Equations*, Academic Press, 1999.
- [4] A. A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, 2006.
- [5] A. Kurt, O. Tasbozan, *Approximate analytical solution of the time fractional Whitham-Broer-Kaup equation using the homotopy analysis method*, Int. J. Pure Appl. Math., **98**(4) (2015), 503-510.
- [6] O. Tasbozan, A. Esen, N. M. Yagmurlu, Y. Ucar, *A numerical solution to fractional diffusion equation for force-free case*, Abstr. Appl. Anal., **2013**, Hindawi, (2013).
- [7] C. Celik, M. Duman, *Crank-Nicolson method for the fractional diffusion equation with the Riesz fractional derivative*, J. of Comput. Phys., **231**(4) (2012), 1743-1750.
- [8] Y. Cenesiz, A. Kurt, *New fractional complex transform for conformable fractional partial differential equations*, J. Appl. Math. Stat. Inf., **12**(2) (2016), 41-47.
- [9] T. Abdeljawad, *On conformable fractional calculus*, J. Comput. Appl. Math., **279** (2015), 57-66.
- [10] M. Eslami, H. Rezazadeh, *The first integral method for Wu-Zhang system with conformable time-fractional derivative*, Calcolo, **53**(3) (2016), 475-485.
- [11] H. Aminikhah, A. R. Sheikhan, H. Rezazadeh, *Sub-equation method for the fractional regularized long-wave equations with conformable fractional derivatives*, Sci. Iran. Transaction B, Mech. Eng., **23**(3) (2016), 1048.
- [12] M. S. Osman, A. Korkmaz, H. Rezazadeh, M. Mirzazadeh, M. Eslami, Q. Zhou, *The unified method for conformable time fractional Schrödinger equation with perturbation terms*, Chinese J. Phys., **56**(5) (2018), 2500-2506.
- [13] Y. Cenesiz, D. Baleanu, A. Kurt, O. Tasbozan, *New exact solutions of Burgers' type equations with conformable derivative*, Wave. Random. Complex, **27**(1) (2017), 103-116.
- [14] A. Kurt, O. Tasbozan, D. Baleanu, *New solutions for conformable fractional Nizhnik-Novikov-Veselov system via G'/G expansion method and homotopy analysis methods*, Opt. Quant. Electron., **49**(10) (2017), 333.
- [15] K. Hosseini, P. Mayeli, R. Ansari, *Bright and singular soliton solutions of the conformable time-fractional Klein-Gordon equations with different nonlinearities*, Wave. Random Complex, **28**(3) (2018), 426-434.
- [16] A. Korkmaz, K. Hosseini, *Exact solutions of a nonlinear conformable time-fractional parabolic equation with exponential nonlinearity using reliable methods*, Opt. Quant. Electron., **49**(8) (2017), 278.
- [17] H. Rezazadeh, H. Tariq, M. Eslami, M. Mirzazadeh, Q. Zhou, *New exact solutions of nonlinear conformable time-fractional Phi-4 equation*, Chinese J. Phys., **56**(6) (2018), 2805-2816.
- [18] H. Bulut, T.A. Sulaiman, H.M. Baskonus, H. Rezazadeh, M. Eslami, M. Mirzazadeh, *Optical solitons and other solutions to the conformable space-time fractional Fokas-Lenells equation*, Optik, **172** (2018), 20-27.
- [19] H. Rezazadeh, S. M. Mirhosseini-Alizamini, M. Eslami, M. Mirzazadeh, M. Mirzazadeh, S. Abbagari, *New optical solitons of nonlinear conformable fractional Schrödinger-Hirota equation*, Optik, **172** (2018), 545-553.
- [20] I. E. Inan, *Multiple soliton solutions of some nonlinear partial differential equations*, Univers. J. Math. Appl., **1**(4) (2018), 273-279.
- [21] H. Rezazadeh, M. S. Osman, M. Eslami, M. Ekici, A. Sonmezoglu, M. Asma, W. A. M. Othman, B. R. Wong, M. Mirzazadeh, Q. Zhou, A. Biswas, M. Belic, *Mitigating Internet bottleneck with fractional temporal evolution of optical solitons having quadratic-cubic nonlinearity*, Optik, **164** (2018), 84-92.
- [22] A. Biswas, M. O. Al-Amr, H. Rezazadeh, M. Mirzazadeh, M. Eslami, Q. Zhou, S. P. Moshokoa, M. Belic, *Resonant optical solitons with dual-power law nonlinearity and fractional temporal evolution*, Optik, **165** (2018), 233-239.
- [23] H. Bulut, T. A. Sulaiman, H. M. Baskonus, *Dark, bright optical and other solitons with conformable space-time fractional second-order spatiotemporal dispersion*, Optik, **163** (2018), 1-7.
- [24] M. H. Cherif, D. Ziane, *Homotopy analysis Aboodh transform method for nonlinear system of partial differential Equations*, Univers. J. Math. Appl., **1**(4) (2018), 244-253.
- [25] A. M. Wazwaz, *The extended tanh method for new solitons solutions for many forms of the fifth-order KdV equations*, App. Math. Comput., **184**(2) (2007), 1002-1014.
- [26] D. Ziane, T. M. Elzaki, M. Hamdi Cherif, *Elzaki transform combined with variational iteration method for partial differential equations of fractional order*, Fundam. J. Math. Appl., **1**(1) (2018), 102-108.
- [27] D. Feng, K. Li, *On exact traveling wave solutions for (1+ 1) dimensional Kaup-Kupershmidt equation*, Appl. Math., **2**(6) (2011), 752-756.
- [28] C. A. Gomez S, *New traveling waves solutions to generalized Kaup-Kupershmidt and Ito equations*, Appl. Math. Comput., **216**(1) (2010), 241-250.
- [29] F. Tascan, A. Akbulut, *Construction of exact solutions to partial differential equations with CRE method*, Commun. Adv. Math. Sci., **2**(2) (2019), 105-113.
- [30] M. H. Cherif, D. Ziane, *Variational iteration method combined with new transform to solve fractional partial differential equations*, Univers. J. Math. Appl., **1**(2) (2018), 113-120.
- [31] A. H. Salas, *Solving the generalized Kaup-Kupershmidt equation*, Adv. Studies Theor. Phys., **6**(18) (2012), 879-885.
- [32] R. Khalil, M. Al Horani, A. Yousef, M. Sababheh, *A new definition of fractional derivative*, J. Comput. Appl. Math., **264** (2014), 65-70.
- [33] H. Rezazadeh, A. Korkmaz, M. Eslami, J. Vahidi, R. Asghari, *Traveling wave solution of conformable fractional generalized reaction Duffing model by generalized projective Riccati equation method*, Opt. Quant. Electron., **50**(3) (2018), 150.
- [34] R. Polat, *Finite difference solution to the space-time fractional partial differential-difference Toda lattice equation*, J. Math. Sci. Model., **1**(3) (2018), 202-205.