

## A Note On Simplicial Groupoids

\*Ali Aytekin<sup>1</sup> 

\*<sup>1</sup> Pamukkale University, Faculty of Science and Literature, Department of Mathematics, DENİZLİ

(Alınış / Received: 12.02.2020, Kabul / Accepted: 26.03.2020, Online Yayınlanma / Published Online: 01.04.2020)

### Keywords

Groupoid,  
Simplicial Object,  
Crossed Module

**Abstract:** Showing the equivalence of the two categories is very important in terms of investigating the properties. This equivalence shows that the properties existing in the category exist in the other category without the need to examination. From this point of view, since the category of groupoids is a category better known than the category of simplicial groupoids, thanks to equivalence, all the properties in the category of groupoids are valid for the category of simplicial groupoids. Main objective of this paper is to give simplicial object in category of groupoids and to prove the equivalence between the category of simplicial groupoids and the category of crossed modules of groupoids.

## Simplisel Grupoidler Üzerine Bir Not

### Anahtar Kelimeler

Grupoid,  
Simplisel Obje,  
Çaprazlanmış Modül

**Öz:** İki kategorinin denliğini göstermek özelliklerin incelenmesi bakımında oldukça önemlidir. Bu denklik, kategoride var olan özelliklerin, incelemeye gerek kalmadan diğer kategoride de var olduğunu gösterir. Bu açıdan bakıldığında grupoidler kategorisi, simplisel grupoidler kategorisine göre daha iyi bilinen bir kategori olduğundan denklik sayesinde grupoidler kategorisindeki özelliklerin tamamı simplisel grupoidler kategorisi için geçerlidir. Bu makalenin temel amacı grupoidler kategorisinde simplisel objeyi vermek ve simplisel grupoidler kategorisiyle grupoidler üzerinde çaprazlanmış modüller kategorisinin denliğini göstermektir.

### 1. Introduction

Crossed modules over groups originally were defined by Whitehead [20]. He mentioned crossed modules in this study on which algebraic structures of relative homotopy groups. Since then, notion of crossed module has taken an important place in other areas. Some of studies are [4,5,10,13,14,15] about this topic. Simplicial groups were initially introduced by Kan [17]. After that, different authors defined concept of simplicial such as simplicial Lie algebras, simplicial Leibniz algebras and equivalences in related categories were given in some of the studies [1,2,6,14].

Groupoids were defined by Brandt [7] after a long working process. Afterwards, many studies have been done by researchers. According to Brown [8] groupoids can be considered as the generalization of groups. In this context, it seems possible to transfer the work done on groups to groupoids. See [3,9,18,19] for details about groupoids.

### 2. Material and Method

#### 2.1. Groupoids and Crossed Modules

We recall some definitions from [9].

A groupoid is a small category in which every morphism is an isomorphism, i.e invertible.

That is, there exists morphism  $\alpha^{-1}$  for any morphism  $\alpha$  such that

$$\alpha \circ \alpha^{-1} = e_{t(\alpha)} \text{ and } \alpha^{-1} \circ \alpha = e_{s(\alpha)},$$

where  $e: C_0 \rightarrow C_1$  converts the identity morphism at an object and  $s, t: C_1 \rightarrow C_0$  are group morphisms.

We write a groupoid as  $C = (C_0, C_1)$ , where  $C_0$  is the set of objects and  $C_1$  is the set of morphisms. A groupoid  $C$  is called totally disconnected, if  $C_1(x, y)$  is empty, for all  $x, y \in C_0$  with  $x \neq y$ , i.e  $s = t$ . There exists a morphism of groupoids

$$f = (f_0, f_1): (C_0, C_1) \rightarrow (C'_0, C'_1)$$

such that the diagram

$$\begin{array}{ccc} C_1 & \xrightarrow{f_1} & C'_1 \\ \downarrow s_1 & & \downarrow s'_1 \\ & & \downarrow t'_1 \\ C_0 & \xrightarrow{f_0} & C'_0 \end{array}$$

is commutative. Thus, it can be formed category of groupoids denoted by **Gpd**.

**Examples:**

- i. Every group can be considered as a groupoid with single object.
- ii. A category whose objects are all sets and morphisms are bijective functions is a groupoid.
- iii. A category whose objects are topological spaces and morphisms are homeomorphisms is a groupoid.
- iv. Let  $X$  be a set and  $(x, y)$  be a morphism from  $x$  to  $y$ , for all  $x, y \in X$ . If the composition is defined as

$$(y, z) \circ (x, y) = (x, z)$$

for all  $x, y, z \in X$ , then  $(X, X \times X)$  becomes a groupoid.

- v. Let  $X$  be a set,  $G$  be a group and triple  $(x, g, y)$  be a morphism from  $x$  to  $y$ , for all  $x, y \in X$  and  $g \in G$ . If the composition is defined as

$$(y, h, z) \circ (x, g, y) = (x, hg, z),$$

for all  $g, h \in G$ , then  $(X, X \times G \times X)$  is a groupoid which is called trivial groupoid.

**Remark 2.1.** Let  $f = (f_0, f_1): C \rightarrow C'$  be a morphism of groupoids. Kernel of the morphism is the set  $\ker f = \{a \in C: \text{there exists an } x \in C_0 \text{ such that } f(a) = e_x\}$ .

**Definition 2.2.** Let  $G$  and  $C$  be groupoids over same object set and  $C$  be totally disconnected. Then an action of  $G$  on  $C$  is defined by a function

$$G_s \times_t C \rightarrow C \\ (g, c) \mapsto {}^g c$$

which satisfies

- a)  ${}^g c$  is defined if and only if  $t(c) = s(g)$  and then  $t({}^g c) = t(g)$ ,
- b)  ${}^g(c_1 \circ c_2) = {}^g c_1 \circ {}^g c_2$ ,
- c)  ${}^{g \circ h} c_1 = {}^g({}^h c_1)$  and  $e_x c_1 = c_1$ ,

for all  $c_1, c_2 \in C(x, x)$  and  $g \in G(x, y)$ ,  $h \in G(y, z)$ .

**Definition 2.3.** Let  $G$  and  $C$  be groupoids providing the above conditions. If the morphism  $\partial: C \rightarrow G$  which satisfies

**CM1)**  $\partial({}^g c) = g^{-1} \circ \partial(c) \circ g$

**CM2)**  $\partial(c_1) \circ c = c_1^{-1} \circ c \circ c_1$

for all  $c_1, c \in C(x, x)$ ,  $g \in G(x, y)$ , and  $x, y \in C_0$  then triple  $(C, G, \partial)$  is called a crossed module of groupoids.

A morphism of crossed modules of groupoids from  $(C, G, \partial)$  to  $(C', G', \partial')$  is a pair of groupoids morphisms

$$\alpha: C \rightarrow C' \text{ and } \beta: G \rightarrow G'$$

such that the diagrams

$$\begin{array}{ccc} C & \xrightarrow{\partial} & G \\ \alpha \downarrow & & \downarrow \beta \\ C' & \xrightarrow{\partial'} & G' \end{array} \quad \begin{array}{ccc} G \times C & \longrightarrow & C \\ (\beta, \alpha) \downarrow & & \downarrow \alpha \\ G' \times C' & \longrightarrow & C' \end{array}$$

are commutative. Thus, it can be get category of crossed modules of groupoids denoted by **XMod(Gpd)**.

### 3. Results

We recall some definitions from [12,16].

Let  $\Delta$  be the category of finite ordinals. A simplicial object  $C$  in **Gpd**, or briefly a simp. gpd, is the functor

$$C: \Delta^{op} \rightarrow \mathbf{Gpd}.$$

In other words, let  $C = \{C_0, C_1, \dots, C_n, \dots\}$  be a family of groupoids and

$$d_i^n: C_n \rightarrow C_{n-1} \text{ and } s_i^n: C_n \rightarrow C_{n+1}$$

groupoid morphisms for  $0 \leq i \leq n$ . The triple  $(C, d_i, s_j)$  is called simplicial groupoid if  $d_i$  and  $s_j$  satisfy the following axioms which called simplicial identities

$$\begin{aligned} d_i d_j &= d_{j-1} d_i && \text{for } i < j \\ d_i s_j &= \begin{cases} s_{j-1} d_i & \text{for } i < j \\ id & \text{for } i = j \text{ or } i = j + 1 \\ s_j d_{i-1} & \text{for } i > j + 1 \end{cases} \\ s_i s_j &= s_{j+1} s_i && \text{for } i \leq j, \end{aligned}$$

for  $0 \leq i \leq n$ , where  $d_i$  and  $s_j$  is called face and degeneracy map respectively. Note that face and dejeneracy morphisms of  $C$  are identity on objects. Simplicial groupoids can be pictured as follows;

$$\begin{array}{ccccccc} & & \xrightarrow{\quad} & & \xrightarrow{\quad} & & \xrightarrow{\quad} \\ & & \vdots & & \xrightarrow{d_0, d_1, d_2} & & \xrightarrow{d_0, d_1} \\ C: \dots C_k & \xrightarrow{\quad} & C_{k-1} \dots C_2 & \xrightarrow{\quad} & C_1 & \xrightarrow{\quad} & C_0 \\ & & \xleftarrow{\quad} & & \xleftarrow{s_0, s_1} & & \xleftarrow{s_0} \\ & & \vdots & & & & \\ & & \xleftarrow{\quad} & & & & \end{array}$$

A simplicial groupoid morphism  $f: C \rightarrow C'$  with  $d_i$  and  $s_j$  maps is a family of groupoid homomorphisms  $f_n: C_n \rightarrow C'_n$  such that

$$d_i f_n = f_{n-1} d_i \text{ and } f_n s_i = s_i f_{n-1}$$

for all  $i$  and  $n$ . So, it can be obtained the category of simplicial groupoids denoted by **Simp(Gpd)**.

#### 3.1. The Moore Complex and Truncated Objects

The Moore complex  $NC$  of a simplicial object  $C$  in **Simp(Gpd)** is the sequence

$$NC: \dots \rightarrow NC_n \xrightarrow{\partial_n} NC_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_2} NC_1 \xrightarrow{\partial_1} NC_0$$

where  $NC_0 = C_0$ ,  $NC_n = \bigcap_{i=0}^{n-1} \ker d_i$  and  $\partial_n$  is the restriction of  $d_n$  to  $NC_n$ . It is called that  $NC$  is Moore

complex of simplicial groupoid  $C$  and is of length  $k$  if  $NC_n = 0$ , for all  $n \geq k+1$ . Thus, one can define a category whose objects are simplicial groupoids with Moore complex of length  $k$  and the morphisms are families of homomorphisms compatible with face and degeneracy maps. We denote that category by **Simp** <sub>$\leq k$</sub>  **(Gpd)**.

The following terminology for groupoid case is adapted from [11]. Let  $C_i$  be a groupoid for  $0 \leq i \leq k$ . A simplicial groupoid defined by

$$\begin{array}{ccccccc} & & & \longrightarrow & & & \\ & & & \vdots & & & \\ \dots & \longrightarrow & 0 & \longrightarrow & C_k & \xrightleftharpoons{\quad} & C_{k-1} \dots C_2 & \xrightleftharpoons{\quad} & C_1 & \xrightleftharpoons{\quad} & C_0 \\ & & & \vdots & & & & \xleftarrow{s_0, s_1} & & \xleftarrow{s_0} & \\ & & & \longleftarrow & & & & & & & \end{array}$$

is called  $k$ -truncated simplicial groupoid and denoted by **Tr<sub>k</sub>Simp(Gpd)**. This structure has truncated functor

$$tr_k : \mathbf{Simp}(\mathbf{Gpd}) \rightarrow \mathbf{Tr}_k \mathbf{Simp}(\mathbf{Gpd})$$

has left adjoint  $st_k$  called  $k$ -skeleton and right adjoint  $cost_k$  called  $k$ -coskeleton. These can be shown as follows;

$$\mathbf{Tr}_k \mathbf{Simp}(\mathbf{Gpd}) \xleftarrow[\text{tr}_k]{\text{cost}_k} \mathbf{Simp}(\mathbf{Gpd}) \xleftarrow[\text{tr}_k]{st_k} \mathbf{Tr}_k \mathbf{Simp}(\mathbf{Gpd}).$$

See [11] for details.

**Theorem 3.1** The category **XMod(Gpd)** is naturally equivalent to the category **Simp** <sub>$\leq 1$</sub>  **(Gpd)**.

**Proof:** Let  $C$  be a simplicial groupoid with Moore complex of length 1 and  $C_0, C_1$  be two groupoids. Take  $G = \ker d_0$  and  $\partial$  is the restriction of  $d_1$  to  $G$ . Define the action of  $C_0$  on  $G$  by

$$\begin{aligned} C_0 \times G &\rightarrow G \\ (c, g) &\mapsto {}^c g = s_0(c)^{-1} \circ g \circ s_0(c) \end{aligned}$$

for all  $c \in C_0$  and  $g \in G$ . By using the action given above  $\partial : G \rightarrow C_0$  becomes a crossed module. Because

**CM1)**

$$\begin{aligned} \partial({}^c g) &= \partial(s_0(c)^{-1} \circ g \circ s_0(c)) \\ &= \partial s_0(c)^{-1} \circ \partial(g) \circ \partial s_0(c) \\ &= d_1 s_0(c)^{-1} \circ \partial(g) \circ d_1 s_0(c) \\ &= c^{-1} \circ \partial(g) \circ c \end{aligned}$$

**CM2)**

$$\begin{aligned} \partial({}^{g'} g) &= s_0 \partial(g)^{-1} \circ g' \circ s_0 \partial(g) \\ &= s_0 d_1(g)^{-1} \circ g' \circ s_0 d_1(g) \\ &= s_0 d_1(g)^{-1} \circ g' \circ s_0 d_1(g) \circ \left[ (g^{-1} \circ (g')^{-1} \circ g) \circ (g^{-1} \circ (g') \circ g) \right] \\ &= s_0 d_1(g)^{-1} \circ g' \circ s_0 d_1(g) \circ (g^{-1} \circ (g')^{-1} \circ g) \circ (g^{-1} \circ (g') \circ g) \\ &= d_2 s_0(g)^{-1} \circ d_2 s_1(g') \circ d_2 s_0(g) \circ d_2 s_1(g^{-1}) \circ d_2 s_1(g')^{-1} \circ d_2 s_1(g) \circ (g^{-1} \circ (g') \circ g) \\ &= d_2 (s_0(g)^{-1} \circ s_1(g') \circ s_0(g)) \circ d_2 (s_1(g^{-1}) \circ s_1(g')^{-1} \circ s_1(g)) \circ (g^{-1} \circ (g') \circ g) \\ &= (g^{-1} \circ (g') \circ g) \end{aligned}$$

for all  $c \in C_0$  and  $g, g' \in G$ . So we obtain the functor

$$U : \mathbf{Simp}_{\leq 1}(\mathbf{Gpd}) \longrightarrow \mathbf{XMod}(\mathbf{Gpd}).$$

On the other hand, let  $\partial : G \rightarrow C_0$  be a crossed module. With the action of  $C_0$  on  $G$  one can obtain the semi-direct product

$$G \tilde{\rtimes} C_0 = \{(g, c) : g \in G \text{ and } c \in C_0\}$$

where multiplication of elements is

$$(g_1, c_1)(g_2, c_2) = (g_1 \circ^{c_1} g_2, c_1 \circ c_2)$$

for all  $(g_1, c_1), (g_2, c_2) \in G \tilde{\rtimes} C_0$ . Also, we have the morphism

$$d_0 : G \tilde{\rtimes} C_0 \rightarrow C_0 \\ (g, c) \mapsto c$$

$$d_1 : G \tilde{\rtimes} C_0 \rightarrow C_0 \\ (g, c) \mapsto \partial(g) \circ c$$

$$s_0 : C_0 \rightarrow G \tilde{\rtimes} C_0 \\ c \mapsto (0, c)$$

which satisfy the simplicial identities. Consequently,

$$C_1 \begin{array}{c} \xleftarrow{d_0, d_1} \\ \xrightarrow{s_0} \end{array} C_0$$

is a 1-truncated simplicial groupoid. As a result, we obtain the functor

$$V : \mathbf{XMod}(\mathbf{Gpd}) \longrightarrow \mathbf{Simp}_{\leq 1}(\mathbf{Gpd}).$$

Thus, these functors can be showed as

$$\mathbf{Simp}_{\leq 1}(\mathbf{Gpd}) \begin{array}{c} \xrightarrow{U} \\ \xleftarrow{V} \end{array} \mathbf{XMod}(\mathbf{Gpd}).$$

#### 4. Discussion and Conclusions

In this study, notion of simplicial object was given in the category of groupoids and it was shown that the category of simplicial groupoids is equivalent to the category of crossed modules of groupoids. Thanks to this equivalence, it was understood that these two categories have common properties.

#### References

- [1] Arvasi, Z., Porter, T. 1997. Higher dimensional peiffer elements in simplicial commutative algebras. *Theory and Applications of Categories*, 3(1)(1997), 1-23.
- [2] Ataseven, Ç. 2016. Relations Among Hihger Order Crossed Modules over Groupoids. *Konuralp Journal of Mathematics*, 4(1)(2016), 282-290.
- [3] Atik, M., Aytekin, A., Uslu, E. Ö. 2017. Representability of actions in the category of (Pre)crossed modules in Leibniz algebras. *Communications in Algebra*, 45(5)(2017), 1825-1841.
- [4] Aytekin, A., Casas, J.M., Uslu, E. Ö. 2012. Semi-Complete Crossed Modules of Lie Algebras. *Journal of Algebra and Its Applications*, 11(5)(2012), 1-24.
- [5] Boyacı, Y., Avcıoğlu, O. 2014. Some Relations Between Crossed Modules and Simplicial Objects in Categories of Interest. *European Journal of Pure and Applied Mathematics*, 7(4)(2014), 412-418.
- [6] Brandt, H. 1927. Über eine Verallgemeinerung des Gruppenbegriffes. *Mathematische Annalen*, 96(1)(1927), 360-366.
- [7] Brown, R. 1987. From Groups to Groupoids : a Brief Survey. *Bulletin of the London Mathematical Society*, 19 (2)(1987), 113-134.
- [8] Brown, R., Higgins, P.J., Sivera, R. 2011. *Nonabelian Algebraic Topology: Filtered Spaces, Crossed Complexes, Cubical Homotopy Groupoids*. Zürich, Switzerland: European Mathematical Society Tracts in Mathematics.
- [9] Casas, J.M., Khmaladze, E., Ladra, M. 2008. Crossed Modules for Leibniz nalgebras. *Forum Mathematicum*, 20 (5) 2008, 841-858.

- [10] Curtis, E.B. 1971. Simplicial Homotopy Theory. *Advances in Mathematics*, 6(2)(1971), 107-209.
- [11] Dwyer, W.G., Kan, D.M. 1984. Homotopy Theory and Simplicial Groupoids. *Indagationes Mathematicae*, 46(1984), 379-385.
- [12] Ellis, G.J. 1988. Higher dimensional crossed modules of algebras. *Journal of Pure and Applied Algebra*, 52(3) (1988), 277-282.
- [13] Ellis, G.J. 1993. Homotopical aspects of Lie algebras. *Journal of The Australian Mathematical Society*, 54(3) (1993), 393-419.
- [14] Emir, K., Akay, H.G. 2019. Pullback crossed modules in the category of racks. *Hacettepe Journal of Mathematics and Statistics*, 48(1)(2019), 140-149.
- [15] Garzon, A. R., Miranda, J.G., Osoriok, R. 2000. A Simplicial Description of The Homotopy Category of Simplicial Groupoids. *Theory and Applications of Categories*, 7(14)(2000), 263-283.
- [16] Kan, D.M. 1958. A Combinatorial Definition of Homotopy Groups. *Annals of Mathematics*, 67(2)(1958), 288-312.
- [17] Mucuk, O., Demir, S. 2018. Normality and quotient in crossed modules over groupoids and double groupoids. *Turkish Journal of Mathematics*, 42(5)(2018), 2336-2347.
- [18] Mucuk, O., Kılıcarlan, B., Şahan, T., Alemdar, N. 2011. Group-groupoids and monodromy groupoids. *Topology and its Applications*, 158(2011), 2034-2042.