

Evaluation of Clinical and Radiological Indicators of Childhood Head Trauma

Çocukluk Çağı Kafa Travmalarının Klinik ve Radyolojik Göstergelerinin Değerlendirilmesi

Münever Yılmaz[®], Ayşe Berna Anil[®], Murat Anil[®], Mehmet Helvacı[®]

Cite as: Yılmaz M, Anil AB, Anil M, Helvacı M. Evaluation of clinical and radiological indicators of childhood head trauma. Forbes J Med. 2021;2(2):79-86.

ABSTRACT

Objective: The aim of this study is to determine the clinical signs of traumatic brain injury and its long-term effects on prognosis by evaluating the clinical and radiological findings of the patients admitted to the pediatric emergency department due to blunt head trauma.

Method: The cases who applied to the pediatric emergency department due to head trauma were examined prospectively. Glasgow Coma (GCS) and Pediatric Trauma Scores (PTS) were calculated. The patients were evaluated neurologically 6 months after they were discharged.

Results: A total of 707 pediatric patients [mean age: 59.8 ± 42.6 months; range: 1 month to 13 years; 263 (37.2%) girls] were evaluated prospectively. Pathology was detected in 101 cases (45.9%) [(epidural hematoma, 14; subdural hematoma, 11; brain edema, 36; intracerebral hematoma, 6; subarachnoid hemorrhage, 8; cerebral contusion, 22. Seventy-two (10.1%) patients had skull fractures.] Seventeen cases (2.4%) were operated, and 7 (1.4%) cases were lost. In children aged < 2 years vomiting, tachypnea, focal neurological findings, multitrauma, GCS <15 and low PTS were more common with traumatic brain injury (p <0.05). Vomiting, GCS <15 and low PTS were more common in children >2 years old and with traumatic brain injury (p <0.05). Neurological sequelae were not detected in patients aged <2 years with mild trauma. Loss of consciousness, pulse rate, respiratory and blood pressure abnormalities, focal neurological findings, low GCS and PTS were more common in children aged >2 years and with neurological sequelae (p <0.05).

Conclusion: Physical examination findings, GCS, and PTS levels are useful tools in predicting the short- and long-term consequences of the injury.

Keywords: Pediatric head trauma, Glasgow Coma score, Pediatric Trauma Score, scalp hematoma

ÖZ

Amaç: Bu çalışmanın amacı, künt kafa travması nedeniyle çocuk acil servisine başvuran hastaların yaralanmaya bağlı klinik ve radyolojik bulgularını değerlendirmek; travmatik beyin yaralanmasının klinik işaretlerini ve uzun dönemde prognoza etkilerini belirlemektir.

Yöntem: Çocuk Acil Servisi'ne kafa travması nedeniyle başvuran olgular prospektif olarak incelendi. Olgularda Beyin tomografisi (BT) çekme kararı izleyen hekimin kararına bırakıldı. Olguların Glasgow Koma (GKS) ve Pediatrik Travma Skorları (PTS) hesaplandı. Olgular hastaneden taburcu olduktan 6 ay sonra nörolojik olarak değerlendirildi.

Bulgular: Toplam 707 çocuk hasta [ortalama yaş: 59,8±42,6 ay; dağılım: 1 ay-13 yaş; 263 (%37,2) kız] prospektif olarak değerlendirildi. Yüz bir olguda (%45,9) patoloji saptandı [(epidural hematom, 14; subdural hematom, 11; beyin ödemi, 36; intraserebral hematom, 6; subaraknoid kanama, 8; serebral kontüzyon, 22 ve kafa kırığı, 72 (%10,1)]. On yedi (2,4%) hasta operasyona alındı. Yedi (%1,4) olgu kaybedildi. İki yaşından küçüklerde kusma, takipne, fokal nörolojik bulgu, multitrauma, GKS<15 ve düşük PTS saptananlarda travmatik beyin yaralanması daha sıkı (p<0,05). İki yaşından büyüklerde kusma, GKS<15 ve düşük PTS saptananlarda travmatik beyin yaralanması daha sıkı (p<0,05). Altıncı ayda iki yaşından küçük minör tıvmalı hastalarda nörolojik sekel saptanmadı. İki yaşından büyüklerde travma sonrası bilinç kaybı öyküsü, nabız sayısı, solunum ve kan basıncı anormallikleri, fokal nörolojik bulgu, düşük GKS ve PTS nörolojik sekel kalanlarda daha sıkı (p<0,05).

Sonuç: Fizik bakı bulguları, GKS ve PTS düzeyleri yaralanmanın kısa ve uzun dönem sonuçlarını öngörmeye faydalı araçlardır.

Anahtar Kelimeler: Pediyatrik kafa travması, Glasgow Koma skoru, Pediatrik Travma Skoru, skalp hematomu

Received/Geliş: 14.12.2020

Accepted/Kabul: 26.02.2021

Publication online: 13.08.2021

Sorumlu Yazar/

Corresponding Author:

Münever Yılmaz

Department of Pediatrics, Pamukkale
University Medical Faculty,
Denizli, Turkey

✉ munevveryl@yahoo.com

ORCID: 0000-0002-5076-1479

A.B. Anil 0000-0003-3670-3771

Department of Pediatrics, Izmir Katip
Celebi University Medical Faculty,
Izmir Turkey

M. Anil 0000-0002-2596-4944

Department of Pediatrics, Izmir
Democracy University Medical
Faculty, Izmir, Turkey

M. Helvacı 0000-0003-3265-8475

Department of Pediatrics, Health
Sciences University, Izmir Tepecik
Teaching and Training Hospital, Izmir



INTRODUCTION

Traumas are one of the most important causes of child death in the world.¹ In every 100,000 children, 1,100 children are admitted to the emergency room for head trauma, while 31 of them are hospitalized and 3.4 of them die.² Most common causes of head trauma include falls from heights, and traffic accidents.³

Most of the children presented with head trauma are cases with minor head trauma and a small number of children need surgery.³ Computed tomography (CT) is an effective and practical method of showing traumatic brain injury in children. However, radiation exposed during CT, can cause cancers in the long term. Therefore, doctors should decide whether or not to perform CT in pediatric patients based on the physical examination findings, age, and cause of the head trauma.^{3,4}

The aim of this study was to evaluate the clinical and radiological findings of patients admitted to the pediatric emergency department due to blunt head trauma and to determine the clinical signs of traumatic brain injury and its effects on prognosis in the long term.

MATERIALS AND METHODS

Between 01.9.2005-31.8.2006, patients under the age of 14 who applied to the Pediatric Emergency Department for head trauma were evaluated prospectively. The patients who were referred to another hospital or refused treatment, and those with missing emergency records were excluded from the study. At admission, Glasgow Coma (GCS) and Pediatric Trauma Scores (PTS) of the cases were calculated.⁵ Those with GCS 13-15 were evaluated as "minor head trauma".⁶ The scalp hematomas of the cases were divided into four groups according to their size as nonexistent, barely palpable, easily palpable and large scalp hematomas.⁷ In patients aged 24 months or less, the fall distance of 1 meter, and in patients older than 2 years the fall distance of ≥ 1.5 meters were accepted as the criteria for falling from height.³ Falling from a distance less than these

heights was considered as simple fall.

The decision to obtain CT was made by the pediatrician who was in duty in the emergency department. No standard algorithm was used. Radiologists evaluated the CTs. The neurosurgeons evaluated the cases for surgical indications. All discharged children were evaluated by telephone calls within the 48th hour, on the 7th day, at the 1st month after trauma. Six months after the trauma, the cases were evaluated neurologically. The cases were divided into ≤ 24 , and >24 month- old groups according to their age. The results were analyzed with SPSS 15.0 (SPSS for Windows 15.0, IL, USA). Chi-square, Student-t, One-way ANOVA, Mann-Whitney U and Kruskal-Wallis tests were used. $P < 0.05$ was considered statistically significant.

This study was approved by Tepecik Hospital Education and Scientific Commission as a thesis study.

RESULTS

A total of 707 pediatric patients [mean age: 59.8 ± 42.6 months; median age: 53 months; range: 1 month-13 years; 263 (37.2%) girls and 444 (62.8%) boys] were evaluated prospectively. When the patients were divided into 2 groups as aged < 2 (192 patients, 27.2%) and >2 (515 patients 72.8%) years, their mean ages were 13.8 ± 6.3 and 76.9 ± 37.3 months, respectively. The highest number of admissions was in July (n: 130 ; 18.4%). The most common cause of trauma was simple fall (n: 309; 43.7%) (Table 1).

The median GCS of the patients at admission was 15 points (interquartile range: 12-15). According to the firstly estimated GCS, 665 (94%) patients had mild, 21 (3%) had moderate and 21 (3%) had severe head traumas. The median PTS of the cases was 12 (interquartile range: 7-12). In a total of 100 patients (14.1%) multiple organ injuries [musculoskeletal injury in 78, (11%); cervical injury in 10 (1.4%); abdominal injury in 9 (1.3%); and genitourinary injury in 3 (0.04%) cases] were detected.

Table 1. Etiologic distribution of pediatric head traumas by months

Months n (%)	Simple fall N: 309 n (%)	Falling from a height N: 119 n (%)	Traffic accident N: 29 n (%)	Pedestrian accident N: 120 n (%)	Bicycle accident N: 55 n (%)	Assault N: 7 n (%)	Other N: 68 n (%)
January	11 (42.3)	4 (15.4)	1 (3.8)	3 (11.6)	0	0	7 (26.9)
February	16 (52.2)	2 (6.9)	2 (6.9)	3 (10.3)	0	1 (3.4)	5 (17.2)
March	27 (62.8)	7 (16.3)	1 (2.3)	5 (11.6)	0	2 (2.3)	2 (4.7)
April	12 (25)	13 (27.1)	5 10.4	10 (20.8)	3 (6.3)	1 (2.1)	4 (8.3)
May	18 (30.5)	11 (18.6)	3 (5.1)	24 (40.7)	1 (1.7)	0	2 (3.4)
June	34 (31.8)	21 (19.6)	2 (1.9)	30 (28)	14 (13.1)	0	6 (5.6)
July	67 (51.5)	22 (16.9)	4 (3.1)	11 (8.)	14 (10.8)	0	12 (9.2)
August	36 (37.9)	15 (15.8)	7 (7.4)	19 (20)	10 (10.5)	1 (1.1)	7 (7.4)
September	55 (53.4)	19 (18.4)	2 (1.9)	3 (2.9)	13 (12.6)	1 (0.9)	10 (9.8)
October	12 (50)	3 (12.5)	2 (8.3)	5 (20.8)	0	0	2 (8.3)
November	11 (47.6)	1 (4.8)	0	3 (14.2)	0	0	7 (33.3)
December	11 (50)	1 (4.5)	0	4 (18.2)	0	2 (9.1)	4 (18.2)




Figure 1. Clinical results of children with blunt head trauma

A total of 220 patients (31.1%) underwent CT. Traumatic injury was detected in 101 cases (45.9% of the cases were identified with CT) [epidural hematoma in 14 (1.9%); subdural hematoma in 11 (1.6%); brain edema in 36 (5.1%); intracerebral hematoma (0.8%), subarachnoid hemorrhage in 8 (1.1%) and cerebral contusion in 22 (3.1%) cases]. Seventy-two (10.1%) patients had cranial fractures [linear fractures in 63 (8.9%); depressed fractures in 6 (0.8%) and basilar skull fractures in 3 (0.4%) cases]. Hundred and twenty-three (17.4%) patients were hospitalized. Seventeen (2.4%) cases were operated. A total of 7 (1.4%) cases exited (Figure 1). When all cases were evaluated at the 6th month after trauma, neurological sequelae were detected in 14 (1.7%) patients including cases with muscle paralysis (n: 8; 1%), strabismus (n:3; 0.4%) , hearing loss (n:1; 0.1%) had, speech impairment (n:1; 0.1%) , vision and hearing loss (n:1 ;0.1%) .

Sixty-five (33.8%) out of the 192 patients aged ≤ 2 years who were admitted to the emergency department due to head trauma had undergone CT. Intracranial injury was detected in 23 (34.5%) patients (Table 2). Among children aged ≤ 2 years, vomiting, abnormal respiratory pattern (tachypnea, bradypnea, abnormal breathing, apnea, etc.), focal neurological findings and multiple injuries were more common in cases with traumatic brain injury ($p < 0.05$). Additionally, GCS and PTS were lower in cases with abnormal CT ($p < 0.05$) (Table 2).

When patients with skull fractures were evaluated, only 6 patients with scalp hematoma were found to have fractures. There was no statistically significant relationship between the size of the scalp hematoma and the presence of fracture ($p > 0.05$). When the relationship between the location of scalp hematoma and the presence of fracture was evaluated, no fractures were detected in any

Table 2. Comparison of the cases with traumatic brain injury or not according to the CT in children ≤ 2 years old

	All patients			Minor head trauma		
	Traumatic brain injury in CT n=23	Normal CT n=42	p	Traumatic brain injury in CT n= 13	Normal CT n= 41	p
Girls, n (%)	10 (43.5)	22 (52.4)	0.670	6 (46.2)	21 (51.2)	0.890
Mean age, month, mean \pm standard deviation	11.5 \pm 7.4	13.2 \pm 5.8	0.800	12.0 \pm 7.9	13.0 \pm 5.7	0.780
Seizure, n (%)	2 (8.2)	5 (11.9)	0.670	2 (15.4)	4 (9.8)	0.778
Vomiting, n (%)	4 (17.4)	21 (50)	<0.001	3 (23.1)	21 (51.2)	0.556
Fever, n (%)	1 (4.3)	1 (2.4)	0.800	0	1 (2.4)	0.560
Pulse abnormality, n (%)	7 (30.4)	5 (1.9)	0.344	0	4 (9.8)	0.110
Respiratory abnormality, n (%)	4 (17.4)	1 (2.4)	0.032	0	1 (2.4)	0.890
Blood pressure abnormality, n (%)	3 (13)	5 (11.9)	0.900	0	4 (9.8)	0.552
Scalp hematoma, n (%)	10 (43.5)	16 (38.1)	0.776	4 (30.8)	16 (39)	0.889
Small scalp hematoma, n (%)	2 (8.2)	3 (7.1)	0.343	1 (7.7)	3 (7.3)	0.900
Medium scalp hematoma, n (%)	3 (13)	6 (14.3)	0.567	1 (7.7)	6 (14.6)	0.098
Large scalp hematoma, n (%)	5 (21.7)	7 (16.7)	0.455	2 (15.4)	7 (17.1)	0.670
Focal neurological sign, n (%)	0	2 (4.8)	<0.001	0	2 (14.9)	0.078
Multiple injuries, n (%)	5 (21.7)	1 (2.4)	<0.001	1 (7.7)	0	0.340
GCS, median (IQR)	11 (8-15)	15 (13-15)	<0.001	15 (14-15)	15 (14-15)	0.890
PTS, median (IQR)	7 (3-11)	10 (8-12)	<0.001	11 (10-12)	10 (10-12)	0.894

CT: Computed tomography; GCS: Glasgow Coma Score PTS: Pediatric Trauma Score; IQR: interquartile range

of the patients with frontal and occipital hematoma, while linear fractures were detected in 2 of the 11 patients with parietal scalp, and in 2 of the 4 patients with temporal scalp hematomas, while basilar fracture were detected in 1 patient ($p<0.001$).

Hundred and eighty-one out of 192 (94.3%) patients aged ≤ 2 years who were admitted to the emergency department due to head trauma had minor head trauma. In this group, 54 (29.8%) cases had CT. Traumatic injuries (skull fractures in 12, subdural hematoma in 1, and cerebral edema in 1 case) were detected in 13 (24.1%) cases. In this group, no parameters were significant compared to pathologic findings (if any) detected in CT ($P>0.05$). Three (1.7%) patients with minor head trauma were operated. Neurological sequelae were not detected in any of the cases with minor head trauma (Table 2).

Seven (3.6%) out of the 192 patients aged ≤ 2 years old who were admitted to the emergency department due to head trauma were operated. The mean age (14.1 ± 6.3 months vs 8.7 ± 6.2 months; $p= 0.03$), GCS (11 vs 15; $p<0,001$) and PTS (8 vs 10; $p<0.001$) were comparatively lower in patients aged ≤ 2 years. Four children (2.1%) had neurological sequelae at the end of the 6th month. Focal neurological signs and multiple injuries were more common ($p<0.05$); GCS and PTS were lower ($p<0.05$) in cases with neurological sequelae (Table 3).

Hundred and fifty-five (29.5%) out of 515 patients aged >2 years who were admitted to the emergency department due to head trauma had undergone CT. Traumatic injury was detected in 77 (49.7%) patients. Vomiting and focal neurological signs were more common in those with abnormal CT group. GCS and PTS were lower in patients with traumatic brain injury ($p<0.05$) (Table 4).

Table 3. Comparison of children ≤ 2 years old in terms of surgical treatment and development of neurological sequelae

	Surgical treatment (+) n: 7	Surgical treatment (-) n: 185	p	Neurological sequelae (+) n= 4	Neurological sequelae (-) n= 188	p
Girl, n (%)	1 (14.2)	81 (43.8)	0.065	1 (25)	81 (43.1)	0.780
Mean age, month, mean \pm standard deviation	8.7 ± 6.2	14.1 ± 6.3	0.030	15.2 ± 4.8	13.8 ± 6.4	0.087
Seizure, n (%)	0	7 (3.9)	0.654	0	7 (3.7)	0.590
Vomiting, n (%)	1 (14.2)	30 (16.2)	0.890	0	31 (16.5)	0.060
Pulse abnormality, n (%)	2 (28.6)	19 (10.3)	0.344	3 (75)	18 (9.5)	0.051
Respiratory rate abnormality, n (%)	1 (14.2)	5 (2.7)	0.760	2 (50)	4 (2.1)	0.055
Blood pressure abnormality, n (%)	1 (14.2)	11 (5.9)	0.560	2 (50)	10 (5.3)	0.780
Scalp hematoma, n (%)	2 (28.6)	54 (29)	0.060	2 (50)	54 (28.7)	0.340
Small scalp hematoma, n (%)	0	23 (12.4)	0.450	1 (25)	22 (11.7)	0.600
Middle scalp hematoma, n (%)	2 (28.6)	18 (9.7)	0.235	1 (25)	19 (10.1)	0.560
Large scalp hematoma, n (%)	0	13 (7)	0.110	0	13 (6.9)	0.070
Focal neurological sign, n (%)	0	37 (20)	0.340	2 (50)	4 (2.1)	<0.001
Multiple injuries, n (%)	0	6 (3.2)	0.980	2 (50)	11 (5.9)	<0.001
GCS, median (IQR)	11 (9-14)	15 (13-15)	<0.001	7 (5-9)	15 (13-15)	<0.001
PTS, median (IQR)	8 (5-9)	10 (8-12)	<0.001	1 (5-9)	10 (8-12)	<0.001

CT: Computed tomography; GCS: Glasgow Coma Score PTS: Pediatric Trauma Score; IQR: interquartile range

Table 4. Comparison of the cases with head trauma in children >2 years old

	Traumatic brain injury in CT n= 78	Normal CT n= 77	P	Operation (+) n: 10	Operation (-) n: 505	p	Neurological sequelae (+) n: 10	Neurological sequelae (-) n: 505	p
Female gender, n (%)	32 (41)	27 (35.1)	0.440	3 (30)	178 (35.2)	0.567	3 (30)	178 (35.2)	0.456
Loss of consciousness, n (%)	26 (33.3)	25 (32.5)	0.901	5 (50)	50 (9.9)	<0.001	4 (40)	51 (10.1)	<0.001
Seizure, n (%)	7 (8.9)	7 (10.9)	0.980	1 (10)	13 (2.6)	0.785	1 (10)	13 (2.6)	0.233
Vomiting, n (%)	32 (41)	44 (57.1)	0.040	3 (30)	87 (17.2)	0.125	2 (20)	88 (17.4)	0.098
Fever, n (%)	0	2 (2.6)	0.152	1 (10)	11 (2.2)	0.343	3 (30)	10 (1.9)	<0.001
Pulse abnormality, n (%)	7 (8.9)	2 (2.6)	0.095	1 (10)	29 (5.7)	0.451	2 (20)	27 (5.3)	<0.001
Respiratory abnormality, n (%)	12 (15.4)	8 (10.4)	0.356	1 (10)	31 (6.1)	0.123	3 (30)	29 (5.7)	<0.001
Blood pressure abnormality, n (%)	9 (11.5)	10 (12.9)	0.780	5 (50)	141 (27.9)	0.065	2 (20)	144 (28.5)	0.065
Small size scalp hematoma, n (%)	28 (35.9)	25 (32.5)	0.650	2 (20)	57 (11.3)	0.554	0 (0)	59 (11.7)	0.655
Medium size scalp hematoma, n (%)	8 (10.2)	7 (10.9)	0.960	0	63 (12.5)	0.041	1 (10)	62 (12.3)	0.076
Large size scalp hematoma, n (%)	9 (11.5)	11 (14.3)	0.370	3 (30)	20 (3.9)	0.022	1 (10)	22 (4.4)	0.230
Focal neurological sign, n (%)	8 (10.2)	2 (2.6)	0.041	5 (50)	50 (9.9)	<0.001	4 (40)	51 (10.1)	<0.001
Multiple traumas, n (%)	19 (24.4)	12 (15.6)	0.175	1 (10)	13 (2.6)	0.760	1 (10)	13 (2.6)	0.330
Age, month, mean± standard deviation	74.5 ± 35.6	73.4 ± 38.8	0.721	89.7± 50.7	76.6±37	0.066	74.4±26.8	76.9± 7.5	0.890
GCS, median (IQR)	12 (10-14)	14 (11-15)	<0.001	11 (10-13)	15 (13-15)	<0.001	11 (10-14)	15 (13-15)	<0.001
PTS, median (IQR)	10 (3-9)	12 (11-12)	<0.001	9 (7-10)	11 (10-12)	<0.001	4 (6-12)	12 (11-12)	<0.001

CT: Computed tomography; GCS: Glasgow Coma Score PTS: Pediatric Trauma Score; IQR: interquartile range

Ten (1.9%) cases older than two years were operated. Post-traumatic loss of consciousness ($p<0.001$), medium and large scalp hematoma ($p= 0.04$ and $p=0.02$) were more common in children undergoing operation. GCS ($p<0.001$) and PTS ($p=0.001$) were lower in patients who needed surgery (Table 4).

Neurological sequelae were detected in 10 (1.9%) of 515 patients aged >2 years after 6 months. Posttraumatic loss of consciousness ($p<0.001$), abnormal pulse rate ($p<0.001$), abnormal blood pressure ($p<0.001$), respiratory abnormality ($p<0.001$), fever ($p<0.001$), and existence of focal neurological finding ($p<0.001$) were more

common, while GCS ($p < 0.001$) and PTS ($p < 0.001$) were lower.

Four hundred and eighty-two (93.6%) out of 525 patients aged > 2 years had mild head trauma. Of these, 123 (25.5%) underwent CT and 52 (42.3%) had traumatic brain injury. Only PTS was significantly lower in patients with mild head trauma who had traumatic brain injury ($p = 0.01$). Of these patients, 5 (1%) were operated and 4 (0.8%) had neurological sequelae.

DISCUSSION

In our study, more than 700 pediatric head traumas were evaluated prospectively. Nearly three-quarters of cases were over two years old, and the most frequent hospital admission occurred in July. The most common cause of head trauma was simple falls. Traumatic brain injury was detected in 14.3% of all pediatric head trauma cases. Of these, 2.4% were operated and mortality was less than 1%. Neurological sequelae were detected in 1.9% of survivors. The presence of vomiting, abnormality in vital findings, focal neurological findings, multiple traumas, low GCS and PTS levels, middle / large size scalp hematoma, parietal and/or temporal scalp hematomas were important predictors for clinical outcomes.

The vast majority of head traumas in children is due to falls from a height followed by traffic and bicycle accidents.^{1,2} The results of our study were similar with the literature. In more than half of the cases, the cause of head trauma was fall from a height, followed by traffic and bicycle accidents. Due to the characteristics of the summer season, the pediatric traumas occurred mostly in July.

Traumatic brain injury in CT is detected between 3-5% of children under two years of age with minor head traumas. Clinically significant head trauma requires surgery in 1% and 0.2% of the cases.^{3,8} The incidence of traumatic brain injury in cases aged > 2 years with minor head traumas ranges between 3-7%. Clinically significant head trauma requires surgery in 1% and 0.1-0.6% of the cases.^{3,8,9} When GCS decreased, the risk of

traumatic brain injury increases.^{8,9} In our study, the rates of brain injury and the need of surgery were about twice of the literature. Less than 1% of the cases deceased. About 2% of the children had neurological sequelae. The different results obtained may be associated with the characteristics of the countries in which the studies were conducted. Publications in the literature are usually carried out in the developed western countries. The hospital where our study was conducted serves to the lowest socioeconomic region of Izmir, Turkey's third largest city. This may be the reason why traumas are more frequent and more severe.

Change of consciousness in the emergency department, loss of consciousness after trauma, headache and vomiting are common findings in children presenting with head trauma.^{10,11} Clinical signs indicating abnormal level of consciousness such as agitation, lethargy, and $GCS < 15$ are the strongest indicators of traumatic brain injury.³ One of the most serious pathological consequences of head trauma is increased intracranial pressure. Vomiting, headache, low GCS, focal neurological findings, abnormalities in vital findings are important clinical indicators for increased intracranial pressure.¹² In our study, vomiting, respiratory abnormality, focal neurological findings, low levels of GCS and PTS were more common in traumatic brain injuries. Similarly, other than vomiting, the same findings were more frequent in patients with neurological sequelae. In other words, clinical signs pointing to pathology in CT also predict long-term prognosis.

Especially in infants, large scalp hematoma, especially hematoma of the temporal or parietal regions is considered one of the important indicators of traumatic brain injury.¹³ In our study, the temporal or parietal scalp hematomas in children ≤ 2 years old were more often accompanied by cranial fracture. However, no risk factors related to hematoma size were detected in this age group. In those over two years old, the large size of the hematoma was an indicator of a skull fracture.

In children affected by blunt high-energy trauma, PTS, measured in the emergency department, is a good marker in predicting the need for critical intervention and the risk of mortality. However, considering all traumas, it has a moderate effectiveness in predicting hospitalization.¹⁴ PTS expresses the level of consciousness, age of the patient, skeletal injury level and airway patency by one number.⁵ In our study, PTS was found as an effective scoring system in predicting traumatic brain injury, the risk for surgery and neurological sequelae in all age groups.

In conclusion, although head trauma is common in children, the risk of related death and neurological sequelae is generally low. Physical examination findings, GCS and PTS levels are useful tools for predicting short- and long-term consequences of pediatric blunt head traumas.

Ethics Committee Approval: This study was approved by Tepecik Hospital Education and Scientific Commission as a thesis study.

Conflict of Interest: None.

Funding: None.

Informed Consent: Informed consent form was received from the participant parents.

REFERENCES

1. <http://www.cdc.gov/ncipc/wisqars> (Accessed on July 14, 2020).
2. Surveillance report of traumatic brain injury-related emergency department visits, hospitalizations, and deaths. Centers for Disease Control and Prevention, 2014. https://www.cdc.gov/traumaticbraininjury/pdf/TBI-Surveillance-Report-FINAL_508.pdf (accessed April 22, 2020).
3. Kuppermann N, Holmes JF, Dayan PS, et al.: Pediatric Emergency Care Applied Research Network (PECARN). Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. *Lancet*. 2009;374(9696):1160-70. doi: 10.1016/S0140-6736(09)61558-0
4. Bressan S, Romanato S, Mion T, Zanconato S, Da Dalt L. Implementation of adapted PECARN decision rule for children with inör head injury in the pediatric emergency department. *Acad Emerg Med*. 2012;19(7):801-7. doi: 10.1111/j.1553-2712.2012.01384.x.
5. Furnival RA, Schunk JE. ABCs of scoring systems for pediatric trauma. *Pediatr Emerg Care*. 1999;15(3):215-23. doi:10.1097/00006565-199906000-00013
6. Carroll LJ, Cassidy JD, Holm L, Kraus J, Coronado VG; WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. Methodological issues and research recommendations for mild traumatic brain injury: the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med*. 2004;(43 Suppl):113-25. doi: 10.1080/16501960410023877.
7. Bin SS, Schutzman SA, Greenes DS. Validation of a clinical score to predict skull fracture in head-injured infants. *Pediatr Emerg Care*. 2010;26(9):633-9. doi: 10.1097/PEC.0b013e3181ef0440
8. Osmond MH, Klassen TP, Wells GA, et al; Pediatric Emergency Research Canada (PERC) Head Injury Study Group. CATCH: a clinical decision rule minor use of computed tomography in children with minor head injury. *CMAJ*. 2010;182(4):341-8. doi: 10.1503/cmaj.091421
9. Da Dalt L, Marchi AG, Laudizi L et al. Predictors of intracranial injuries in children after blunt head trauma. *Eur J Pediatr*. 2006;165(3):142-8. doi: 10.1007/s00431-005-0019-6
10. Pandor A, Goodacre S, Harnan S, et al. Diagnostic management strategies for adults and children with minor head injury: a systematic review and an economic evaluation. *Health Technol Assess*. 2011;15(27):1-202. doi: 10.3310/hta15270.
11. Lee LK, Monroe D, Bachman MC, et al. Isolated loss of consciousness in children with minor blunt head trauma. *JAMA. Pediatr* 2014; 168:837-47. doi: 10.1001/jamapediatrics.2014.361.
12. Miller Ferguson N, Shein SL, Kochanek PM, et al. Intracranial Hypertension and Cerebral Hypoperfusion in Children with Severe Traumatic Brain Injury: Thresholds and Burden in Accidental and Abusive Insults. *Pediatr Crit Care Med*. 2016;17(5):444-50. doi: 10.1097/PCC.0000000000000709.
13. Schutzman SA, Nigrovic LE, Mannix R. The Infant Scalp Score: A Validated Tool to Stratify Risk of Traumatic Brain Injury in Infants With Isolated Scalp Hematoma. *Acad Emerg Med*. 2020. doi: 10.1111/acem.14087.
14. Anil M, Saritaş S, Bıçlıoğlu Y, Gökalp G, Kamit Can F, Anil AB. Pediyatrik Travma Skorunun Çocuk Acil Servisindeki Performansı: İleriye Yönelik Bir Çalışma. *J Pediatr Emerg Intensive Care Med*. 2017;4:1-7. doi: 10.4274/cayd.30074