

# YAYIN BİLGİSİ

A Young Boy with Renal Agenesis and Ectopic Seminal Vesicle: Zinner Syndrome

Ahmet Yasin Yitik, Furkan Ufuk, Sinan Çelen

Journal of Urological Surgery - 2023

Zinner syndrome is a rare disorder, and it is associated with unilateral renal agenesis, ipsilateral ejaculatory duct atresia (obstruction), and cystic dilatation of the ipsilateral seminal vesicle. Here, we present an eighteen-year-old male with Zinner syndrome who presented to a urology clinic with perineal/pelvic pain and dysuria. We also emphasize the computed tomography and magnetic resonance imaging findings of Zinner syndrome.

Anahtar kelimeler: Imaging, congenital malformations, renal agenesis



10.4274/jus.galenos.2022.2022.0015



R9EPGYsBQzmg-9NMh1Sy



0

Atıf



3

Görüntülenme



1

İndirilme



0

Dinlenme

Copyright © Sobiad Atıf Dizini - [atif.sobiad.com](http://atif.sobiad.com)

# A Young Boy with Renal Agenesis and Ectopic Seminal Vesicle: Zinner Syndrome

Ahmet Yasin Yitik<sup>1</sup>, Furkan Ufuk<sup>1</sup>, Sinan Çelen<sup>2</sup>

<sup>1</sup>Pamukkale University Faculty of Medicine, Department of Radiology, Denizli, Türkiye

<sup>2</sup>Pamukkale University Faculty of Medicine, Department of Urology, Denizli, Türkiye

## Abstract

Zinner syndrome is a rare disorder, and it is associated with unilateral renal agenesis, ipsilateral ejaculatory duct atresia (obstruction), and cystic dilatation of the ipsilateral seminal vesicle. Here, we present an eighteen-year-old male with Zinner syndrome who presented to a urology clinic with perineal/pelvic pain and dysuria. We also emphasize the computed tomography and magnetic resonance imaging findings of Zinner syndrome.

**Keywords:** Imaging, congenital malformations, renal agenesis

## Introduction

Zinner syndrome is a rare disease that is seen in males. It occurs due to a developmental abnormality of the distal part of the Wolffian duct. Zinner first described the Zinner syndrome in 1914 as unilateral renal agenesis, ipsilateral ejaculatory duct obstruction, and ipsilateral seminal vesicle cyst (1-3). It is also accepted as the male equivalent of Müllerian agenesis (Mayer-Rokitansky syndrome) seen in females. Zinner syndrome is associated with unilateral renal agenesis, ipsilateral ejaculatory duct atresia (obstruction), and cystic dilatation of the ipsilateral seminal vesicle (3). The pathogenesis of Zinner syndrome is related to the mesonephric (Wolffian) duct (4). The mesonephric (Wolffian) duct develops into the prostatic part of the urethra, ductus deferens, seminal vesicle, and epididymis owing to testosterone and anti-müllerian hormone (5). An abnormality of development of the distal part of the Wolffian duct can be because by atresia (obstruction) of the ejaculatory duct, leading to the dilatation and cysts of the ipsilateral seminal vesicle. Furthermore, the concurrent ureteral budding abnormalities can lead to renal agenesis or dysplasia. These developmental pathologies can result in azoospermia and oligospermia, which cause primer infertility in males (3-5). Also, seminal vesicle cysts and cystic dilatation of the ipsilateral seminal vesicle can

have mass effects and pelvic perineal pain (4,5). Diagnosis of Zinner syndrome can be made by ultrasonography or computed tomography (CT) scan; but magnetic resonance imaging (MRI) gives a better resolution of the local anatomy. Treatment is usually conservative, but surgery may be required in selected cases.

## Case Report

An eighteen-year-old boy presented to the urologic clinic with pelvic pain and dysuria. Physical examination and laboratory test results, including complete biochemical profile, C-reactive protein, blood count, and urinalysis results, were within normal limits. A urine specimen for culture was obtained. An abdominal ultrasound showed right renal agenesis and a 31x40x52 mm heterogeneous cystic lesion in the right lower quadrant. Unenhanced abdominal CT showed the right renal agenesis and ipsilateral pelvic lesion with well-defined borders (Figure 1). The following pelvic MRI revealed that the ipsilateral pelvic lesion was a cyst with proteinaceous/hemorrhagic content and was compatible with an ectopic and dilated right seminal vesicle (Figure 2). Zinner syndrome should be considered first in the differential diagnosis, characterized by unilateral renal agenesis, ipsilateral ejaculatory duct atresia (obstruction), and cystic

**Correspondence:** Ahmet Yasin Yitik MD, Pamukkale University Faculty of Medicine, Department of Radiology, Denizli, Türkiye

**Phone:** +90 258 296 60 00 **E-mail:** ahmetyasinyitik@gmail.com **ORCID-ID:** orcid.org/0000-0002-1793-7911

**Received:** 08.02.2022 **Accepted:** 07.05.2022

**Cite this article as:** Yitik AY, Ufuk F, Çelen S. A Young Boy with Renal Agenesis and Ectopic Seminal Vesicle: Zinner Syndrome. J Urol Surg, 2023;10(1):74-76.

©Copyright 2023 by the Association of Urological Surgery / Journal of Urological Surgery published by Galenos Publishing House.



dilatation of the ipsilateral seminal vesicle (1). After diagnosis, sperm analysis results were within normal limits, except for mild hemospermia. Laparoscopic surgical resection of the right seminal vesicle was performed due to hemospermia and clinical symptoms in this case. The patient was discharged without complications on the second day after the operation. The one-year follow-up of the patient was uneventful and her clinical complaints wholly regressed. Informed consent was obtained from the patient.

## Discussion


Zinner syndrome is an uncommon anomaly, and its frequency is not known precisely. According to a study on ultrasonography, the incidence of seminal vesicle cysts in patients with ipsilateral renal dysplasia or aplasia (agenesis) is 0.0046% (6). The first-line examination of the assessment of Wolffian duct anomaly is the ultrasound owing to its non-invasive, radiation-free, and easily applicable nature. On ultrasound, seminal cysts are seen as a cystic pelvic mass that can demonstrate thick or irregular wall and mural calcifications. The internal echoes depend on hemorrhage or infection (1,3). Excretory urography and kidney scintigraphy

can indicate ipsilateral kidney agenesis or dysgenesis. Pelvic MRI is an appropriate tool for the detailed examination of pelvic anomalies. It is a gold standard modality for evaluating pelvic abnormalities due to magnificent soft-tissue resolution and multiplanar visualization without radiation. When an absence of urogenital structures is detected on imaging, it should be cautiously examined for possible other Wolffian duct anomalies such as renal agenesis and seminal vesicle cyst (3). While figuring out the etiology of abdominal pain caused by the pressure effects of the seminal vesicle cyst over the adjacent structures, the Zinner syndrome can be diagnosed as in this study (4,5). However, the patients can admit priming infertility caused by azoospermia and oligospermia (6).

Cystic lesions of the seminal vesicle may imitate other pelvic lesions. That's why an accurate diagnosis is required. The differentiation of the cystic pelvic lesions is mainly based on the position as median, paramedian, or lateral (2). Also, abdominal pain causes, such as acute appendicitis, should be excluded. Ultrasonography or CT should be used for differential diagnosis if needed (5,6). Early diagnosis is principal to prevent complications such as infertility in patients with Zinner syndrome and



**Figure 1.** a) Axial unenhanced computed tomography (CT) image at the level of the upper abdomen shows right renal agenesis (arrow). b) Axial CT image at the level of the lower abdomen shows well-defined pelvic lesion (arrowheads). c) Axial CT image at the level of prostate demonstrates the normal left seminal vesicle (arrowhead) and the absence of the right seminal vesicle (arrow)



**Figure 2.** a) T2-weighted and b) T1-weighted coronal magnetic resonance (MR) images of the pelvis show the hyperintense pelvic lesion on both sequences is consistent with a proteinaceous/hemorrhagic cyst (arrowheads). c) T1-weighted axial MR image at the level of prostate demonstrates the normal left seminal vesicle (arrowhead) and the absence of the right seminal vesicle (arrow). The right pelvic mass is consistent with ectopic and a dilated right seminal vesicle, according to pelvic MR imaging findings

avoid nephrotoxic drugs. surgical excision may be considered depending on the size and location of the seminal vesicle cysts and clinical symptoms. Treatment is often conservative if the patient is asymptomatic (1,5). Moreover, since renal agenesis will occur in a patient with Zinner syndrome, a single kidney should be protected by measures such as avoiding nephrotoxic drugs and nephrotoxic contrast agents.

## Conclusion

In conclusion, Zinner syndrome is an unusual anomaly, which includes unilateral kidney agenesis, ipsilateral ejaculatory duct obstruction, and seminal vesicle cyst. Moreover, seminal vesicle ectopia and cystic dilatation may also be seen in patients with Zinner syndrome, as in this case. Zinner syndrome should be kept in mind in the differential diagnosis of young patients with abdominal pain, infertility, and urinary symptoms.

## Ethics

**Informed Consent:** Informed consent was obtained from the patient.

**Peer-review:** Externally peer-reviewed.

## Authorship Contributions

Surgical and Medical Practices: S.Ç., Concept: A.Y.Y., F.U., Design: A.Y.Y., F.U., Data Collection or Processing: A.Y.Y., F.U., S.Ç.,

Analysis or Interpretation: F.U., Literature Search: A.Y.Y., F.U., Writing: A.Y.Y., F.U.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

## References

1. Militaru V, Mihaly ZA, Ilea C, Coman M, Stanciu M, Crisan N, Coman I. Zinner syndrome - case report. *Med Pharm Rep* 2021;94(Suppl No 1):S47-S50.
2. Shebel HM, Farg HM, Kolokythas O, El-Diasty T. Cysts of the lower male genitourinary tract: embryologic and anatomic considerations and differential diagnosis. *Radiographics* 2013;33:1125-1143.
3. Zinner A. Ein fall von intravesikaler Samenblasenzyste. *Wien Med Wochenschr* 1914;64:605-609.
4. Ghonge NP, Aggarwal B, Sahu AK. Zinner syndrome: A unique triad of mesonephric duct abnormalities as an unusual cause of urinary symptoms in late adolescence. *Indian J Urol* 2010;26:444-447.
5. Hannema SE, Hughes IA. Regulation of Wolffian duct development. *Horm Res* 2007;67:142-151.
6. Sheih CP, Hung CS, Wei CF, Lin CY. Cystic dilatations within the pelvis in patients with ipsilateral renal agenesis or dysplasia. *J Urol* 1990;144:324-327.

# BENZER YAYINLAR



The treatment of giant seminal vesicle cyst which accompanying ipsilateral renal agenesis with...  
Bilal Fırat ALP, Zafer DEMİRER, Ali GÜRAĞAÇ, Şeref BAŞAL, Kemal KARA, İbrahim YILDIRIM  
Gülhane Tıp Dergisi - 2015



Radical Surgical Treatment of A Large Seminal Vesicle Cyst Causing Acute Urinary Retention...  
Dejan Djordjevic, Svetomir Dragicevic, Marko Vukovic  
Journal of Urological Surgery - 2021




Pediatric Olguda Alışılmadık Bir Pelvik Ağrı Sebebi: HerlynWernerWunderlich Sendromu  
Yasemin Kayadibi, Elmar Bayraktarov, Evrim Özmen, Altay Gezer, İbrahim Adaletli  
Haseki Tıp Bülteni - 2014



HERLYN-WERNER-WUNDERLICH SYNDROME-ROLE OF IMAGING IN TIMELY AND CORRECT...  
Hidayat Ullah, Atiq Ur Rehman Slehria, Rashid Mahmood, Jehan Akbar  
Pakistan Armed Forces Medical Journal - 2021



Bilateral congenital absence of the vas deferens coexisting with bilateral seminal vesicle...  
Mehmet UMUL, Barış ALTAY, Burak TURNA, Oktay NAZLI  
Süleyman Demirel Üniversitesi Sağlık Bilimleri Dergisi - 2013



Copyright © Sobiad Atıf Dizini - [atif.sobiad.com](http://atif.sobiad.com)