




Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

Poly-arginine conjugated triarylmethyl radical as intracellular spin label



Benoit Driesschaert^{a,b}, Andrey A. Bobko^{a,b}, Timothy D. Eubank^{a,c}, Alexandre Samouilov^d, Valery V. Khrantsov^{a,b}, Jay L. Zweier^{d,*}

^a In Vivo Multifunctional Magnetic Resonance Center, Robert C. Byrd Health Science Center, West Virginia University, Morgantown, WV 26506, United States

^b Department of Biochemistry, West Virginia University School of Medicine, Morgantown, WV 26506, United States

^c Department of Microbiology, Immunology & Cell Biology, West Virginia University School of Medicine, Morgantown, WV 26506, United States

^d Davis Heart & Lung Research Institute, The Ohio State University/Wexner Medical Center, 460 West 12th Avenue, BRT 0390, Columbus, OH 43210, United States

ARTICLE INFO

Article history:

Received 25 January 2016

Revised 16 February 2016

Accepted 17 February 2016

Available online 19 February 2016

Keywords:

EPR

TAM

Poly-arginine

Radical

Cyclic voltammetry

ABSTRACT

Stable triarylmethyl radicals are ideal spin labels used for biomedical electron paramagnetic resonance applications. Previously reported structures exhibit polar charged functions for water solubilization preventing them from crossing the cell membrane. We report the synthesis of a triarylmethyl radical conjugated to poly-arginine peptide allowing intracellular delivery of the paramagnetic label.

© 2016 Elsevier Ltd. All rights reserved.

Stable radicals of tetrathiatriarylmethyl represent soluble spin labels and probes widely used for biomedical electron paramagnetic resonance (EPR) spectroscopy and imaging.¹ They combine excellent biological stability and extraordinary narrow EPR line width as a consequence of their long relaxation times, therefore resulting in a high signal to noise ratio. The six thioacetone moieties provide stability to the radical by electronic and steric effects while the *para*-positions are used to provide water solubility through substitution with polar charged functions such as carboxylic acids (e.g., see Fig. 1 for the structure of **CT-03** derivative also known as Finland trityl). Bimolecular collisions between the paramagnetic probe and molecular oxygen (³O₂) induce a line broadening of the EPR line of the trityl due to pair-wise Heisenberg spin exchange providing a noninvasive, sensitive and quantitative way to measure oxygen in a biological media in vitro and in vivo.^{2,3} Depending on the particular structure of the trityl probe other biological relevant parameters can be measured. For example, the phosphonated derivative allows simultaneous measurement of dissolved oxygen, pH and phosphate concentration in the physiological range in living tissues.^{4,5} Tetrathiatriarylmethyl disulfide biradicals allow for EPR assessment of biologically relevant thiols

such as glutathione (GSH)⁶ while the nitroxide–trityl biradical allows for assessment of tissue redox.⁷

Trityl radicals with charged *para*-functions such as **CT-03** are cell impermeable therefore allowing assessment of important physiological parameters in the extracellular environment.^{5,8} However, intracellular pH, redox status and GSH level provide complementary information of particular importance to physiology and pathology of various diseases such as cancer.⁹ Recently, we synthesized a cell-permeable trityl radical **AMT-02** (Fig. 1) which contain biodegradable acetoxymethyl esters. After crossing the membranes, the esters are hydrolyzed to **CT-03** radical. However, this approach is limited by the low water solubility of **AMT-02**, the slow intra-cellular hydrolysis leading to **CT-03** formation and its subsequent process of excretion from the cells tested.¹⁰ Due to the lipophilic nature of the thioacetone moieties **CT-03** binds to albumin in its sub-domain IIA (Sudlow's site I) and presumably to other biomolecules leading to an increase in the EPR line width. This line broadening leads to a decrease of the signal intensity by a factor of five when excess of bovine serum albumin (BSA) has been used.¹¹ The above factors make application of **AMT-02** probe for the measurement of oxygen complex.¹⁰

Small poly-arginine peptides are known for their ability to cross cell membranes and have been used to deliver nanosized cargo or small molecules inside cells.¹² In this Letter, we explore

* Corresponding author.

E-mail address: jay.zweier@osumc.edu (J.L. Zweier).




Figure 1. Chemical structure of CT-03 and cell permeable AMT-02.

poly-arginine peptides as a carrier for intracellular delivery of a trityl spin label. The activated tri-pentafluorophenol esters **1** (Scheme 1) have been proven to be an excellent intermediate to bind a small peptide to the trityl in a good yield.¹³ Following this strategy, the poly-arginine conjugated trityl **PolyA-TAM** was synthesized. In the first step **CT-03** is activated by oxalyl chloride to lead to the formation of the trityl acyl chloride which is directly engaged in reaction with pentafluorophenol yielding the pentafluorophenol activated ester of **CT-03**. The pure ester is obtained in 75% yield after flash chromatography. Finally coupling the pentafluorophenol ester with commercially available poly-arginine chains ($8 \leq n \leq 12$, MW = 1900 Almenda polymer) in DMSO led to the desired **PolyA-TAM**. After purification by dialysis using 5000 cutoff membrane, the pure radical was obtained in 95% yield as a green solid. The radical was stable, no degradation was observed in water for 24 h at room temperature in oxygenated water.

The EPR spectrum of **PolyA-TAM** in aqueous solution shown in Figure 2. According to the presence of three amide functions, the spectral pattern shows seven partially overlapped lines (1:3:6:7:6:3:1) as a result of hyperfine couplings with three equivalent nitrogen nuclei (^{14}N , $I = 1$) with an hyperfine splitting constant (hfc) of 220 mG, in agreement with the previously reported *para*-amide derivatives of trityl radicals.¹⁴ This spectral assignment supports the presence of three poly-arginine chains bound to the trityl. We anticipated that the charge nature of the poly-arginine chains associated with the bulkiness of our newly synthesized EPR spin label will prevent binding to albumin. In agreement with this hypothesis addition of 600 μM of BSA did not affect EPR spectrum of **PolyA-TAM** except for a small linewidth broadening of 10 mG (See Supplementary data for simulated spectra) apparently due to a change of viscosity of the media.





Figure 2. X-band EPR spectra of 100 μM **PolyA-TAM** solution in deoxygenated 10 mM PBS buffer (pH = 7.4, room temperature) before and after addition of BSA. Acquisition parameters were as follows: sweep width, 3 G; power, 0.2 mW; modulation frequency, 100 kHz; modulation amplitude, 0.05 G; conversion time, 80 ms; time constant, 163.84 ms; resolution, 1024 pt.

Recent study of the metabolism of **CT-03** has shown that the EPR signal can decay due to reduction of the radical under anaerobic conditions¹⁵ by liver microsomes followed by protonation of the anion or by oxidation to cation by some peroxidases or related hemoproteins.¹⁶ The resulting cation can subsequently react with C-, S-, P- or N-nucleophiles.^{17,18} In addition, oxidative decarboxylation of **CT-03** by superoxide radical or peroxy radicals lead to a diamagnetic quinone-methide product.¹⁹ Substitution of the carboxylic acid by amide functions protect against the latter reaction. In order to evaluate the sensitivity toward oxidation and reduction, the redox potentials were measured by cyclic voltammetry. In phosphate buffer, pH = 7.4, the voltammogram shows a reduction of the radical to the anion. The half wave of reduction reaches -0.475 V versus Ag/AgCl indicating that **PolyA-TAM** is easier to reduce than **CT-03** in its carboxylate form (see Table 1). The oxidation of the radical to the cation was observed with a half wave of $E_{1/2} = +0.666$ versus Ag/AgCl indicating that the **PolyA-TAM** is more difficult to oxidize than **CT-03** (Table 1). These results are



Scheme 1. Synthesis of **PolyA-TAM** from **CT-03**.

Table 1
Oxidation and reduction half wave potentials for **PolyA-TAM** and **CT-03**

Compound	Solvent	$E_{1/2}$ ox ^a	$E_{1/2}$ red ^a
CT-03 ¹³	PBS pH = 7.4	+0.434	−0.642
PolyA-TAM	Water pH = 7.4	+0.666	−0.475

^a Calculated according to $E_{1/2} = (E_{pa} + E_{pc})/2$.




Figure 3. Cyclic voltammogram of **PolyA-TAM** recorded in water at pH = 7.4, scan rate = 0.2 V/s.




Figure 4. X-band EPR spectra of internalized **PolyA-TAM** measured after different incubation times at 37 °C. Acquisition parameters were as following: sweep width, 10 G; power, 0.63 mW; modulation frequency, 100 kHz; modulation amplitude, 0.1 G; conversion time, 80 ms; time constant, 367.28 ms; resolution, 1024 pt.

consistent with a higher electron withdrawing effect of amide function compared with the carboxylate ones (Fig. 3).

To investigate the ability of **PolyA-TAM** probe to permeate the cellular membrane and accumulate in intracellular space, MDA-MB-231 cells were incubated in the presence of 3 μM of **PolyA-TAM** probe for different times (30 min, 1 h, 2 h). Then cells were washed three times with 20 ml of PBS, trypsinized, collected and centrifuged. Pellet was dispersed in PBS, loaded into glass capillary followed by EPR spectra acquisition. The observed EPR signal intensity (see Fig. 4) corresponds to about 3.5 μM of **PolyA-TAM** in the sample of 7.5×10^7 cell/ml while no EPR signal was found in the supernatant. This corresponds to a 200 μM intracellular concentration of **PolyA-TAM** assuming that the volume of single MDA-MB-231 cell equals is 2000 μm^3 .³ No significant change of the signal intensity was observed after 30 min of incubation indicating that the penetration of the **PolyA-TAM** into the cell is comparatively fast. The total spectral line widths are similar for **PolyA-TAM** measured in buffer and inside the cells (cf. Figs. 2 and 4). The disappearance of additional hyperfine splitting in cellular sample might arise from broadening of individual components due to higher intracellular viscosity.

In conclusion, we have synthesized a new trityl spin label linked to three poly-arginine peptides. Due to coupling with three nitrogen nuclei of the amide functions, the spectrum shows a septet pattern limiting its application as oximetric EPR probe and

decreasing the signal to noise ratio. No binding to albumin was observed validating our hypothesis that charged nature of the poly-arginine and the bulkiness of the molecule prevent this interaction. The **PolyA-TAM** is able to penetrate across the cellular membrane and was found to be stable inside the cells for hours. Further improvement of the probe structure can be achieved by introduction of a linker between the parent trityl and the poly-arginine chains therefore eliminating hyperfine splitting-induced line broadening and improving signal-to-noise ratio. This strategy was previously reported to bind **CT-03** to polyethylene glycol moieties through using small deuterated ester linker providing a narrow single line EPR spectrum.²⁰

Acknowledgments

This work was partially supported by NIH grants: EB016096, EB014542, CA194013, CA192064, U54GM104942 and F.S.R Université catholique de Louvain Fellowship awarded to B.D. The WVCTSI is acknowledged for start-up to T.D.U., V.V.K., A.A.B. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmcl.2016.02.048>.

References and notes

- Khramtsov, V. V.; Zweier, J. L. In *Stable Radicals: Fundamentals and Applied Aspects of Odd-Electron Compounds*; Hicks, Robin, Ed.; Wiley & Sons Ltd., 2010; p 537.
- Elas, M.; Ahn, K.-H.; Parasca, A.; Barth, E. D.; Lee, D.; Haney, C.; Halpern, H. J. *Clin. Cancer Res.* **2006**, *12*, 4209.
- Ahmad, R.; Kuppusamy, P. *Chem. Rev.* **2010**, *110*, 3212.
- Dhimitruka, I.; Bobko, A. A.; Eubank, T. D.; Komarov, D. A.; Khramtsov, V. V. *J. Am. Chem. Soc.* **2013**, *135*, 5904.
- Bobko, A. A.; Dhimitruka, I.; Zweier, J. L.; Khramtsov, V. V. *Angew. Chem., Int. Ed.* **2014**, *53*, 2735.
- Liu, Y.; Song, Y.; Rockenbauer, A.; Sun, J.; Hemann, C.; Villamena, F. A.; Zweier, J. L. *J. Org. Chem.* **2011**, *76*, 3853.
- Liu, Y.; Villamena, F. A.; Song, Y.; Sun, J.; Rockenbauer, A.; Zweier, J. L. *J. Org. Chem.* **2010**, *75*, 7796.
- Williams, B. B.; al Hallaq, H.; Chandramouli, G. V. R.; Barth, E. D.; Rivers, J. N.; Lewis, M.; Galtsev, V. E.; Karczmar, G. S.; Halpern, H. J. *Magn. Reson. Med.* **2002**, *47*, 634.
- Khramtsov, V. V.; Gillies, R. J. *Antioxid. Redox Signal.* **2014**, *21*, 723.
- Liu, Y.; Villamena, F. A.; Sun, J.; Wang, T.-Y.; Zweier, J. L. *Free Radical Biol. Med.* **2009**, *46*, 876.
- Song, Y.; Liu, Y.; Liu, W.; Villamena, F. A.; Zweier, J. L. *RSC Adv.* **2014**, *4*, 47649.
- Shin, M. C.; Zhang, J.; Min, K. A.; Lee, K.; Byun, Y.; David, A. E.; He, H.; Yang, V. C. *J. Biomed. Mater. Res. A* **2014**, *102*, 575.
- Driesschaert, B.; Levêque, P.; Gallez, B.; Marchand-Brynaert, J. *Tetrahedron Lett.* **2013**, *54*, 5924.
- Driesschaert, B.; Levêque, P.; Gallez, B.; Marchand-Brynaert, J. *Eur. J. Org. Chem.* **2014**, *2014*, 8077.
- Decroos, C.; Li, Y.; Bertho, G.; Frapart, Y.; Mansuy, D.; Boucher, J.-L. *Chem. Res. Toxicol.* **2009**, *22*, 1342.
- Decroos, C.; Li, Y.; Soltani, A.; Frapart, Y.; Mansuy, D.; Boucher, J.-L. *Arch. Biochem. Biophys.* **2010**, *502*, 74.
- Decroos, C.; Boucher, J.-L.; Mansuy, D.; Xu-Li, Y. *Chem. Res. Toxicol.* **2014**, *27*, 627.
- Decroos, C.; Prange, T.; Mansuy, D.; Boucher, J.-L.; Li, Y. *Chem. Commun.* **2011**, 4805.
- Decroos, C.; Li, Y.; Bertho, G.; Frapart, Y.; Mansuy, D.; Boucher, J.-L. *Chem. Commun.* **2009**, 1416.
- Song, Y.; Liu, Y.; Hemmann, C.; Villamena, F.-A.; Zweier, J.-L. *J. Org. Chem.* **2013**, *78*, 1371.