

# Synthesis of tetracyclic iminosugars fused benzo[e][1,3]thiazin-4-one and their HIV-RT inhibitory activity

Zhuqing Yin<sup>a</sup>, Mo Zhu<sup>a</sup>, Sinan Wei<sup>a</sup>, Jie Shao<sup>a</sup>, Yuheng Hou<sup>a</sup>, Hua Chen<sup>a,b,\*</sup>, Xiaoliu Li<sup>a,b,\*</sup>

<sup>a</sup> Key Laboratory of Chemical Biology of Hebei Province, College of Chemistry and Environmental Science, Hebei University, Baoding 071002, China

<sup>b</sup> National Research Center for Carbohydrate Synthesis, Jiangxi Normal University, Nanchang 330022, China

## ARTICLE INFO

### Article history:

Received 2 November 2015

Revised 20 January 2016

Accepted 18 February 2016

Available online 18 February 2016

### Keywords:

Tetracyclic iminosugar

Benzo[e][1,3]thiazin-4-one

Aza-C-pseudonucleoside

One-pot three-component reaction

Anti-HIV RT activity

## ABSTRACT

Several aza-C-pseudonucleosides bearing 1,3-benzothiazin-4-one (**6** and **7**) were prepared by the one-pot three-component condensation from the iminosugar aldehyde **3**, amino acid ethyl/methyl ester hydrochlorides **4(a–c)**, and 2-mercaptobenzoic acid **5**. After removal of Boc and the isopropylidene groups, the target novel tetracyclic iminosugars fused benzo[e][1,3]thiazin-4-one **1(a–c)** and **2(a–b)** were first afforded by the intramolecular cyclo-amidation reaction. Their structures were determined by their <sup>1</sup>H, <sup>13</sup>C NMR, and HRMS (ESI) spectra and X-ray. The tetracyclic iminosugars **1(a–c)** and **2(a–b)** were examined for their HIV reverse transcriptase (RT) inhibitory activities. The result showed that all compounds could effectively inhibit RT activity. Among them, compound **2a** was the best one with the IC<sub>50</sub> value of RT inhibitory activity of 0.82 μM. Structure–activity relationship analysis suggested that 1'R configuration in the tetracyclic azasugars was of benefit to their anti-HIV RT activity.

© 2016 Elsevier Ltd. All rights reserved.

Iminosugars or azasugars, which exhibit effective inhibition against carbohydrate-processing enzymes, have attracted great interest for their potential clinical applications as anti-HIV, anti-diabetic, and anti-cancer agents and immunomodulators in past decades.<sup>1</sup> The bicyclic azasugars, including the naturally occurring compounds (**A** and **B**)<sup>2</sup> and the synthetic ones (**C** and **D**)<sup>3</sup> (Fig. 1), have also been paid much attention due to their increased possibility leading to discovering new bioactive therapeutic agents. To date, a large number of bicyclic azasugars fused nitrogen heterocycle have been synthesized and evaluated.<sup>4</sup> Much less efforts have been put into multicyclic analogs.<sup>5</sup> The tetracyclic azasugars were so far scarcely reported for their synthesis and biological activity study.


Recently, a series of novel bi- and tricyclic thiazolidin-4-one- and benzothiazin-4-one-fused azasugars (**E** and **F**, Fig. 1) have been found to exhibit strong HIV reverse transcriptase (HIV-RT) inhibitory activities.<sup>6</sup> It was well known that most HIV-1 nonnucleoside reverse transcriptase inhibitors (NNRTIs) generally showed butterfly-like conformation when they were binding with HIV-RT.<sup>7</sup> Taking account this dominant conformation as determinant for anti-HIV activity, many families of NNRTIs were initially designed, including the butterfly-like conformationally constrained NNRTIs **G** (MEN 10979) and **H** (Fig. 1).<sup>8</sup> Inspired by this, we conceived that the tetracyclic azasugars fused benzo[e][1,3]thiazin-4-one expanded from the tricyclic ones might be more beneficial for their

binding with HIV-RT due to their possible butterfly-like conformation, like compounds **G** and **H**. Therefore, in this Letter, we would like to first report the design and the synthesis of the novel tetracyclic azasugars fused 1,3-benzothiazin-4-one (**1** and **2**, Fig. 2) as a continuation of our researches. Such newly synthetic azasugars were evaluated for their HIV-RT inhibitory activity in order to further investigate the structure–activity relationship (SAR).


The synthesis of the target tetracyclic azasugars was achieved in three steps. The key reaction for the intermediate aza-C-pseudonucleosides bearing 1,3-benzothiazin-4-one (**6** and **7**) was the one-pot three-component condensation from the iminosugar aldehyde **3**,<sup>9</sup> amino acid ethyl/methyl ester hydrochlorides **4(a–c)** (neutralized by NaHCO<sub>3</sub>), and 2-mercaptobenzoic acid **5** at 40–60 °C as shown in Scheme 1. In the presence of the condensation reagent *N,N'*-dicyclohexyl-carbodiimide (DCC) and the promoter 4-dimethylamino-pyridine (DMAP), the one-pot synthesis afforded the diastereomeric products **6** and **7** (Table 1) in the overall yields of 12.4–59.7% following our reported procedure.<sup>10</sup> However, the reactions were accompanied with the generation of the byproducts **10** and **11** which were directly condensed from the aldehyde **3** and 2-mercaptobenzoic acid **5**. The consumption of the aldehyde **3** maybe caused the low yields of the three-component reactions. Moreover, steric hindrance from the methyl on L- and D-alanine might be another reason for the low yields. Especially in entry 3 (Table 1) using D-alanine methyl ester hydrochloride **4c** as amine source, **6c** was just obtained in 12.4% yield. Although only one isomer was found in this case, we could

\* Corresponding authors. Tel./fax: +86 312 5971116.

E-mail addresses: [hua-todd@163.com](mailto:hua-todd@163.com) (H. Chen), [lixl@hbu.cn](mailto:lixl@hbu.cn) (X. Li).



**Figure 1.** The structures of some bi/tricyclic azasugars **A–F** and conformationally constrained NNRTIs **G** and **H**.




**Figure 2.** The synthetic tetracyclic azasugars fused benzo[e][1,3]thiazin-4-one.

not absolutely rule out small amounts of the other product **7c** from the reaction, its presence was not evident from examination of the crude reaction mixtures. After Boc and the isopropylidene groups

in **6** and **7** were removed in 90%  $CF_3COOH$  to achieve the corresponding products **8** and **9**, then the treatment of **8** or **9** with triethylamine at 50 °C gave the targeted tetracyclic azasugars fused benzo[e][1,3]thiazin-4-one **1** and **2** in good yields by intramolecular cycloamidation from secondary amine and ester. However, it should be mentioned that, under the same condition, the final cyclization could not be effectively performed using  $\beta$ -alanine as amine source which would form seven-membered ring.

The structures of all the newly synthesized tetracyclic iminosugars were determined by their  $^1H$ ,  $^{13}C$  NMR, and HRMS (ESI) spectra. Both analytical and spectral data of compounds are in agreement with the proposed structures. The typical coupling



**Scheme 1.** The synthesis of the tetracyclic azasugars **1(a–c)** and **2(a–b)** using iminosugar aldehyde **3** as starting material. Reagents and conditions: (I) toluene,  $NaHCO_3$ , DCC, DMAP, 40–60 °C; (II) 90%  $CF_3COOH-H_2O$ , rt; (III) MeOH,  $NEt_3$ , 50 °C.

**Table 1**

The synthesis of aza-C-pseudonucleosides bearing 1,3-benzothiazin-4-one (**6** and **7**) by the one-pot three-component condensation

Entry	Amino acid ester hydrochloride	Products		Total yields <sup>a</sup> (%)	Ratio of <b>6</b> / <b>7</b> <sup>b</sup>
		<b>6</b> -1'S	<b>7</b> -1'R		
1	Glycine	30.2	29.5	59.7	1:1
2	L-Alanine	22.1	14.7	36.8	1.5:1
3	D-Alanine	12.4	— <sup>c</sup>	12.4	1:0

<sup>a</sup> Isolated yield.

<sup>b</sup> Determined by <sup>1</sup>H NMR.

<sup>c</sup> Not found.

constants of 1-H and 1'-H (Table 2) indicated that compounds **1(a–c)** with small  $J_{1,1'}$  values should have *cis*-relationship between 1-H and 1'-H, while their corresponding diastereomers are of *trans* forms. Thus, the absolute configuration of compounds **1(a–c)** could be determined to be of (1'S), the others were of (1'R). The absolute configuration of compound **2a** was determined to be of (1'R) by its X-ray crystallographic data (Fig. 3),<sup>11</sup> being consistent with the above NMR results.


HIV-1 reverse transcriptase (RT) inhibitory activity was preliminarily evaluated with the tetracyclic azasugars **1** and **2** (Fig. 2) by determining their percentage inhibition of HIV-RT activity in HIV-1-RT kit by comparison with AZT.<sup>12</sup> The results are shown in Table 3. It could be seen that all the newly synthesized tetracyclic azasugars **1** and **2** showed significant HIV-RT inhibitory activity, better than that of positive control AZT. Especially, compounds **2a** showed a more significant HIV-RT inhibitory activity with the IC<sub>50</sub> value of 0.82 μM which indicated that **2a** might be better accommodated into the HIV-1 RT binding site. The inhibitory activities of the tetracyclic azasugars **2(a–b)** with 1'R configuration are much higher than those of their stereoisomers **1(a–b)** with 1'S configuration, respectively, which indicated that the stereochemistry of C-1' in *R* configuration would be favorable to the HIV-RT inhibitory activities of the compounds.

In conclusion, novel tetracyclic azasugars fused benzo[e][1,3]thiazin-4-one **1** and **2** were first synthesized by the intramolecular cyclo-amidation reaction of the unprotected aza-C-pseudonucleosides, which were afforded by the one-pot three-component

**Table 2**

The coupling constants (Hz) of 1-H and 1'-H in **1** and **2**

<i>cis</i> (1'S)	$J_{1,1'}$ /Hz	<i>trans</i> (1'R)	$J_{1,1'}$ /Hz
<b>1a</b>	4.8	<b>2a</b>	7.2
<b>1b</b>	3.6	<b>2b</b>	10.2
<b>1c</b>	3.6		



**Figure 3.** X-ray crystallographic structure of **2a**.

**Table 3**

In vitro HIV-1-RT kit assay for the tetracyclic azasugars

Compds	IC <sub>50</sub> (μM) (HIV-RT kit assay)	Compds	IC <sub>50</sub> (μM) (HIV-RT kit assay)
<b>1a</b>	8.21 ± 1.24	<b>2a</b>	0.82 ± 0.27
<b>1b</b>	9.65 ± 2.36	<b>2b</b>	4.21 ± 1.03
<b>1c</b>	2.01 ± 0.68	AZT	20.14 ± 1.32

condensation from the iminosugar aldehyde, amino acid ester hydrochlorides, and 2-mercaptobenzoic acid. Their structures were determined by NMR, HRMS (ESI) spectra and X-ray crystallographic data. All the newly synthesized tetracyclic azasugars showed significant HIV-RT inhibitory activity. Among them, compound **2a** was the best one with the IC<sub>50</sub> value of RT inhibitory activity of 0.82 μM. The structure activity relationship (SAR) analysis indicated that the stereochemistry of C-1' in *R* configuration would be favorable to their anti-HIV-RT inhibitory activity.

### Acknowledgments

The financial supports from the National Natural Science Foundation of China (NSFC) (21372060), Hebei Province Natural Science Fund for Distinguished Young Scholars (B2015201005), and the Open Project Program of National Research Center for Carbohydrate Synthesis, Jiangxi Normal University (GJDTZX-KF-2014).

### Supplementary data

Supplementary data (experimental procedures and characterization data for compounds **1(a–c)** and **2(a–b)**) associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmcl.2016.02.049>.

### References and notes

- (a) Compain, P.; Martin, O. R. *Iminosugars: From Synthesis to Therapeutic Applications*; John Wiley & Sons Ltd: Chichester, England, 2007; (b) Asano, N. In *Glycoscience: Chemistry and Chemical Biology* In Fraser-Reid, B. O., Tatsuta, K., Thiem, J., Eds.; Springer-Verlag: Berlin, 2008; pp 1887–1911; (c) Ye, X. S.; Sun, F.; Liu, M.; Li, Q.; Wang, Y. H.; Zhang, G. S.; Zhang, L. H.; Zhang, X. L. *J. Med. Chem.* **2005**, *48*, 3688; (d) Vyavahare, V. P.; Chakraborty, C.; Maity, B.; Chattopadhyay, S.; Puranik, V. G.; Dhavale, D. D. *J. Med. Chem.* **2007**, *50*, 5519.
- (a) Elbein, A. D.; Tropea, J. E.; Mitchell, M.; Kaushal, G. P. *J. Biol. Chem.* **1990**, *265*, 15599; (b) Aoyagi, T.; Suda, H.; Uotani, K.; Kojima, F.; Aoyama, T.; Horiguchi, K.; Hamada, M.; Takeuchi, T. *J. Antibiot.* **1992**, *45*, 1404.
- (a) Benlifa, M.; Isabel García-Moreno, M.; Mellet, C. O.; García-Fernández, J. M.; Wadouachi, A. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 2805; (b) Zou, W.; Bhasin, M.; Vembaiyan, K.; Williams, D. T. *Carbohydr. Res.* **2009**, *344*, 1024; (c) Moncayo, M. A.; Mellet, C. O.; García-Fernández, J. M.; Isabel García-Moreno, M. *J. Org. Chem.* **2009**, *74*, 3595; (d) Enderlin, G.; Taillefumier, C.; Didierjean, C.; Chapleur, Y. *J. Org. Chem.* **2009**, *74*, 8388; (e) Ben-Othman, R.; Othman, M.; Ciamala, K.; Knorr, M.; Strohmman, C.; Decroix, B. *Tetrahedron* **2009**, *65*, 4846; (f) Yadav, L. D. S.; Awasthi, C. *Carbohydr. Res.* **2010**, *345*, 318; (g) Ritthiwigrom, T.; Willis, A. C.; Pyne, S. G. *J. Org. Chem.* **2010**, *75*, 815; (h) Chen, M. J.; Tsai, Y. M. *Tetrahedron* **2011**, *67*, 1564; (i) Bowen, E. G.; Wardrop, D. J. *Org. Lett.* **2010**, *12*, 5330.
- (a) Stütz, A. *Iminosugars as Glycosidase Inhibitors: Nojirimycin and Beyond*; Wiley-VCH: Weinheim, 1999; (b) Afarinkia, K.; Bahar, A. *Tetrahedron: Asymmetry* **2005**, *16*, 1239; (c) Pearson, M. S. M.; Mathe- Allainmat, M.; Fargeas, V.; Lebreton, J. *Eur. J. Org. Chem.* **2005**, *11*, 2159.
- Nair, V.; Purdy, D. F. *Tetrahedron Lett.* **1991**, *32*, 7503.
- (a) Li, X. L.; Qin, Z. B.; Yang, T. Y.; Zhang, H. Z.; Wei, S. N.; Li, C. X.; Chen, H.; Meng, M. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 2712; (b) Chen, H.; Yang, T. Y.; Wei, S. N.; Zhang, H. Z.; Li, R.; Qin, Z. B.; Li, X. L. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 7041; (c) Chen, H.; Li, R.; Liu, Z. Y.; Wei, S. N.; Zhang, H. Z.; Li, X. L. *Carbohydr. Res.* **2013**, *365*, 1; (d) Chen, H.; Hao, L.; Zhu, M.; Yang, T. Y.; Wei, S. N.; Qin, Z. B.; Zhang, P. Z.; Li, X. L. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 3426.
- (a) Zhan, P.; Liu, X. Y.; Li, Z. *Curr. Med. Chem.* **2009**, *16*, 2867; (b) de Béthune, M. P. *Antiviral Res.* **2010**, *85*, 75; (c) Prajapati, D. G.; Ramajayam, R.; Yadav, M. R.; Giridhar, R. *Bioorg. Med. Chem.* **2009**, *17*, 5744; (d) Schäfer, W.; Friebe, W. G.; Leinert, H.; Mertens, A.; Poll, T.; von der Saal, W.; Zilch, H.; Nuber, B.; Ziegler, M. L. *J. Med. Chem.* **1993**, *36*, 726.
- (a) Zhan, P.; Chen, X. W.; Li, D. Y.; Fang, Z. J.; De Clercq, E.; Liu, X. Y. *Med. Res. Rev.* **2013**, *33*, E1; (b) Bellarosa, D.; Antonelli, G.; Bambacioni, F.; Giannotti, D.; Viti, G.; Nannicini, R.; Giachetti, A.; Dianzani, F.; Witvrouw, M.; Pauwels, R.

- Desmyter, J.; De Clercq, E. *Antiviral Res.* **1996**, *30*, 109; (c) Silvestri, R.; Artico, M.; Bruno, B.; Massa, S.; Novellino, E.; Greco, G.; Marongiu, M. E.; Pani, A.; De Montis, A.; La Colla, P. *Antiviral Chem. Chemother.* **1998**, *9*, 139.
9. Li, X. L.; Wei, S. N.; Zhang, H. B.; Zhang, P. Z.; Wang, W.; Chen, H. *Chin. J. Org. Chem.* **2012**, *32*, 1708.
10. (a) Chen, H.; Jiao, L. L.; Guo, Z. H.; Li, X. L.; Ba, C. L.; Zhang, J. C. *Carbohydr. Res.* **2008**, *343*, 3015; (b) Chen, H.; Yin, Q. M.; Li, C. X.; Wang, E. K.; Gao, F.; Zhang, H. B.; Yin, Z.; Wei, S. N.; Li, X. L.; Meng, M.; Zhang, P. Z.; Li, N.; Zhang, J. C. *ACS Med. Chem. Lett.* **2011**, *2*, 845.
11. CCDC-1433230 for **2a** contain the supplementary crystallographic data for this Letter. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
12. Reverse Transcriptase Assay, Colorimetric kit, Roche Diagnostics GmbH, Roche Applied Science, Sandhofer Strasse 116, D-68305 Mannheim, Germany.